

# The New Minimal Supersymmetric GUT

Charanjit S. Aulakh and Sumit K. Garg

*Dept. of Physics, Panjab University  
Chandigarh, India 160014  
E-mail: aulakh@pu.ac.in*

## ABSTRACT

The Supersymmetric SO(10) GUT based on the  $\mathbf{210} \oplus \mathbf{10} \oplus \mathbf{120} \oplus \mathbf{126} \oplus \overline{\mathbf{126}}$  Higgs system and only spontaneous CP violation has 24 real superpotential parameters. It is compatible with all current low energy data and yet has enough freedom to accommodate the still unknown Leptonic CP violation and neutrino mass scale parameters. It is thus the New Minimal Supersymmetric Grand Unified Theory(NMSGUT). We compute the superheavy mass spectrum, couplings and associated gauge evolution threshold effects and determine the regions of the parameter space compatible with perturbative unification, baryon stability and Weinberg angle while permitting spontaneous CKM CP violation. We show that the structural requirements of spontaneous CP violation always tend to raise the unification scale  $M_X$  and with it the mass of all superheavy multiplets and thus to suppress Baryon decay. These scales can be raised even close to the Planck scale so that the NMSGUT gauge Landau pole becomes innocuous. This scenario lends support to our earlier speculations regarding GUT dynamical symmetry breaking and induced gravity via UV condensation at this Landau pole. Fermion mass formulae and the effective superpotential for Baryon violating  $d = 5$  processes - which contains novel channels due to the  $\mathbf{120}$  Higgs - are derived to prepare for the next round of detailed analysis of the phenomenology of this model.

# 1 Introduction

The discovery of neutrino mass was both preceded by[1] and itself provoked[2, 3, 4, 5, 6, 7] intensive investigation of unifying theories that naturally incorporate supersymmetry and the seesaw mechanisms[8] : in particular models with the Left-Right gauge group as a part of the gauge symmetry and  $B - L$  broken at a high scale and R/M-parity preserved to low energies[1]. The close contiguity of the seesaw scale and the Grand Unified scale pushed SO(10) GUTs, which are the natural GUT home of both Type I and Type II seesaw mechanisms, but were long relegated as baroque cousins of the -seemingly- more elegant minimal SU(5) GUT, into centre stage. The understanding that the Susy SO(10) based on the  $\mathbf{210} \oplus \mathbf{10} \oplus \mathbf{126} \oplus \overline{\mathbf{126}}$  Higgs system proposed[10, 11] long ago was the best candidate for the Minimal Supersymmetric GUT (MSGUT) crystallized after the demonstration of its minimality on parameter counting grounds and an elegant reduction of its spontaneous symmetry breaking problem to a single cubic equation with just one unknown parameter[12]. Careful computations of the symmetry breaking and mass spectra [10, 11, 12] became available[3, 4, 5, 6] MSGUT. These theories naturally maintain a structural distinction between Higgs and matter fields and therefore naturally preserve R-parity down to low energies[13, 1, 2].

The initial euphoria[7] that the version utilizing only  $\mathbf{10}, \overline{\mathbf{126}}$  Higgs representations might prove sufficient[9] to fit all low energy fermion data in an elegant and predictive way ran aground when the successful generic fits of fermion mass data were shown to be unrealizable[14, 15] in the context of the actual Seesaw mechanisms available in the MSGUT. Thereafter the GUT based on the  $\mathbf{210} \oplus \mathbf{10} \oplus \mathbf{120} \oplus \mathbf{126} \oplus \overline{\mathbf{126}}$  Higgs system has emerged[14, 15, 16, 17, 18] as the New Minimal Supersymmetric GUT (NMSGUT) capable of fitting all the known fermion mass and mixing data. Remarkably this compatibility has been demonstrated even when all the superpotential parameters are real and with 3 matter yukawas set to zero and the  $\overline{\mathbf{126}}$  vev fixed in an ad-hoc way[18]. Thus at least 4 free parameters are still available to fit the still unknown values of the leptonic CP violation phases, neutrino mass scale and  $\theta_{13}^P$  when they become available. The theory with the  $\mathbf{120}$  thus escapes the contradiction with the composition of the light Higgs doublets which was demonstrated[14, 15] to occur in the MSGUT without it. Thus this actual or New MSGUT calls for and deserves the same detailed analysis of its superheavy threshold effects, fermion mass fit compatibility and exotic effect effective superpotential that we earlier provided for the theory without the  $\mathbf{120}$  [10, 3, 5, 14, 15] which we previously called the MSGUT[12] but whose claim on that name is now tenuous and faded.

Even after it fits the fermion data the NMSGUT must still face the challenge posed by the non observation of Baryon decay in spite of the lore that minimal Susy GUTs imply baryon decay rates via  $d = 5$  operators which are higher than the current experimental upper bounds[21]. Moreover all the analysis of MSGUTs

has so far been done in the framework of softly broken global supersymmetry with no attempt to incorporate its dynamical or spontaneous breaking. Furthermore the NMSGUT suffers[22] from a Landau pole at  $\Lambda_X$  lying just above the perturbative unification scale  $M_X$  due to the huge  $SO(10)$  gauge beta functions implied by the large representations used.

Last week, as this paper was being finalized, a very interesting paper[23] appeared which gave a lucid discussion of these essential and perennial issues[24] and suggested that they could be addressed simultaneously by considering metastable Susy breaking vacua in the context of  $N=1$  Supergravity with non-renormalizable operators included in the matter superpotential to allow escape of the unification scale and  $d = 5$  baryon decay amplitudes from the tight confines[21] of the minimal  $SU(5)$  SGUT. The authors argued that the MSSM singlets of the MSGUT itself could play the role of the supersymmetry breaking field(s) of the Polonyi or other hidden sector and that this melded optimally with a  $M_X \sim M_{Planck}$  solution to the  $d = 5$  baryon decay rate difficulty of Susy GUTs. For reasons we shall explain below we do not find a non-renormalizable superpotential cogent or attractive inasmuch as it puts paid to the predictivity engendered by renormalizability. It seems moreover that it has been invoked more to provide additional parameters that may be fine tuned to satisfy the needs of the attractive phenomenological scenario suggested[23]. It is natural to enquire whether the same program might be naturally fulfilled in a realistic theory with only a renormalizable superpotential such as the MSGUT for which we had computed the threshold corrections already[5, 14, 15] and the NMSGUT which we were in the process of reporting. In fact such a raising of the unification scale was observed by us in considerable regions of the parameter space already[5, 14, 15]. Since however the focus had been on raising neutrino masses by lowering the right handed neutrino mass we had not yet questioned whether the new found[16, 17] leeway in fitting neutrino masses via Type I seesaw with highly suppressed  $\overline{126}$  yukawa couplings[15, 16, 17] would allow us to exploit the domination of the raised unification scale regions in the viable parameter space of the MSGUT and NMSGUT. We therefore remedied this defect and report on the very encouraging indications in detail below. We note that a novel type of “intermediate” scale Dirac supermultiplet supermultiplet transforming as  $[8, 2, \pm 1]$  with respect to the SM gauge group - which couples to quarks via the  $\overline{126}$  yukawa - becomes lighter as the unification scale is raised. It is the effects of this multiplet that will alter the RG equations sufficiently to raise  $\Delta_G$  consistently. Unlike similar multiplets used in[23] for the same purpose, however, the mass of this multiplet does not seem to fall below  $10^{15} GeV$  and that too only when  $M_X$  and  $\alpha_G$  have long become unphysical.

We have speculated[25] that the vice of Asymptotic Strength(AS) of (N)MSGUTs may be turned to good account to construct a theory of dynamical calculable GUT symmetry breaking using an extension of the techniques for dealing with strongly coupled supersymmetric theories[26], and even that AS Susy GUTs(ASSGUTs), which

determine by their RG flow their own physical UV cutoffs, escape the objections to the induced gravity program that halted its development in the 80's[27]. This induced gravity program would gain plausibility if the Planck scale and  $\Lambda_X$  coincided. Such a coincidence is an obvious consequence of raising the perturbative unification scale to just below the Planck scale. We shall see that the requirement of viable unification in the presence of high scale threshold corrections in the NMSGUT leads us almost inevitably to regions of the parameter space where  $M_X$ , together with baryon decay mediating triplet masses, are raised above  $10^{16} GeV$  and  $\Lambda_X$  approaches closer to  $M_{Planck}$ .

The computation of mass spectra, couplings and gauge threshold effects in the NMSGUT and a partial analysis of their implications is the subject of this paper. A calculation of the chiral mass spectra using different conventions and method is already available [31]. However, since our methods also yield all couplings which we need for an evaluation of the phenomenological implications, we present all results consistently using our own conventions.

In Section 2 we recapitulate our notation and the basics of such models. Detailed accounts of our techniques have already been given earlier[3, 5, 14]. In Section 3 and Appendix A we give the mass spectra and in Appendix B we describe an  $SU(5)$  reassembly crosscheck of the spectra we obtain. In Section 4 we discuss how the threshold effects calculated using these spectra determine regions where perturbative unification is viable and sub-regions of these where Baryon decay is naturally suppressed. We shall see how both in the general case of complex superpotential parameters and when the CKM CP violation is spontaneous one is generically led, in the latter case inescapably, towards a raised unification scale: thus leading to a potential resolution of the basic problems of Susy GUTs discussed above, without any contrived cancellations. In Section 5 we give the fermion mass formulae in the presence of the **120** plet using analytic expressions for the null eigenvectors(after fine tuning to keep one pair of doublets light) of the  $6 \times 6$  Higgs doublet  $([1, 2, \pm 1])$  mass matrix (Appendix C). In Section 6 we integrate out the heavy triplets that mediate Baryon decay via  $d = 5$  operators and give the resultant effective superpotential in terms of the matter superfields of the effective MSSM. We conclude with a brief discussion of issues and on-going/future work in Section 7.

## 2 The New Minimal Susy GUT

### 2.1 MSGUT couplings, vevs and masses

The original MSGUT [10, 11, 9, 12] was the renormalizable globally supersymmetric  $SO(10)$  GUT whose Higgs chiral supermultiplets consist of AM type totally antisymmetric tensors : **210**( $\Phi_{ijkl}$ ), **126**( $\bar{\Sigma}_{ijklm}$ ), **126**( $\Sigma_{ijklm}$ )( $i, j = 1...10$ ) which break the GUT symmetry to the MSSM, together with Fermion mass (FM) Higgs **10**-plet(**H<sub>i</sub>**).

The  $\overline{126}$  plays a dual or AM-FM role since it also enables the generation of realistic charged fermion and neutrino masses and mixings (via the Type I and/or Type II Seesaw mechanisms); three **16**-plets  $\Psi_A (A = 1, 2, 3)$  contain the matter including the three conjugate neutrinos ( $\bar{\nu}_L^A$ ).

The superpotential (see[12, 3, 4, 5] for comprehensive details ) contains the mass parameters

$$m : \mathbf{210}^2 \quad ; \quad M : \mathbf{126} \cdot \overline{\mathbf{126}}; \quad M_H : \mathbf{10}^2 \quad (1)$$

and trilinear couplings

$$\lambda : \mathbf{210}^3 \quad ; \quad \eta : \mathbf{210} \cdot \mathbf{126} \cdot \overline{\mathbf{126}}; \quad \gamma \oplus \bar{\gamma} : \mathbf{10} \cdot \mathbf{210} \cdot (\mathbf{126} \oplus \overline{\mathbf{126}}) \quad (2)$$

In addition one has two complex symmetric matrices  $h_{AB}, f_{AB}$  of Yukawa couplings of the the  $\mathbf{10}, \overline{\mathbf{126}}$  Higgs multiplets to the **16.16** matter bilinears. The  $U(3)$  ambiguity due to  $SO(10)$  ‘flavour’ redefinitions can be used to remove 9 of the 24 real parameters in  $f, h$ . In addition rephasing of the remaining 4 fields  $\Phi, \mathbf{H}, \Sigma, \bar{\Sigma}$  removes 4 phases from the 14 parameters in  $m, M, M_H, \lambda, \eta, \gamma, \bar{\gamma}$  leaving 25 superpotential parameters to begin with. Strictly speaking, since a fine tuning to keep one pair of doublets light is an intrinsic part of the MSGUT scenario, an additional parameter (say  $M_H$  ) may be considered as fixed so that there are actually 24 free superpotential parameters. In addition the electroweak scale vev  $v_W$ , and  $\tan \beta$  are relevant external parameters for the light fermion spectrum determined by the GUT yukawa structures.

The GUT scale vevs that break the gauge symmetry down to the SM symmetry (in the notation of[3]) are[10, 11]

$$\langle (15, 1, 1) \rangle_{210} : \langle \phi_{abcd} \rangle = \frac{a}{2} \epsilon_{abcdef} \epsilon_{ef} \quad (3)$$

$$\langle (15, 1, 3) \rangle_{210} : \langle \phi_{ab\tilde{\alpha}\tilde{\beta}} \rangle = \omega \epsilon_{ab} \epsilon_{\tilde{\alpha}\tilde{\beta}} \quad (4)$$

$$\langle (1, 1, 1) \rangle_{210} : \langle \phi_{\tilde{\alpha}\tilde{\beta}\tilde{\gamma}\tilde{\delta}} \rangle = p \epsilon_{\tilde{\alpha}\tilde{\beta}\tilde{\gamma}\tilde{\delta}} \quad (5)$$

$$\langle (10, 1, 3) \rangle_{\overline{126}} : \langle \bar{\Sigma}_{\hat{1}\hat{3}\hat{5}\hat{8}\hat{0}} \rangle = \bar{\sigma} \quad (6)$$

$$\langle (\overline{10}, 1, 3) \rangle_{126} : \langle \Sigma_{\hat{2}\hat{4}\hat{6}\hat{7}\hat{9}} \rangle = \sigma. \quad (7)$$

The vanishing of the D-terms of the  $SO(10)$  gauge sector potential imposes only the condition  $|\sigma| = |\bar{\sigma}|$ . Except for the simpler cases corresponding to enhanced unbroken gauge symmetry ( $SU(5) \times U(1), SU(5), G_{3,2,2,B-L}, G_{3,2,R,B-L}$  etc)[12, 4], this system of equations is essentially cubic and can be reduced to a single equation [12] for a variable  $x = -\lambda\omega/m$ , in terms of which the vevs  $a, \omega, p, \sigma, \bar{\sigma}$  are specified :

$$C(x, \xi) = 8x^3 - 15x^2 + 14x - 3 + \xi(1 - x)^2 = 0 \quad (8)$$

where  $\xi = \frac{\lambda M}{\eta m}$ . Then the dimensionless vevs in units of  $(m/\lambda)$  are  $\tilde{\omega} = -x$  [12] and

$$\tilde{a} = \frac{(x^2 + 2x - 1)}{(1 - x)} \quad ; \quad \tilde{p} = \frac{x(5x^2 - 1)}{(1 - x)^2} \quad ; \quad \tilde{\sigma}\tilde{\bar{\sigma}} = \frac{2\lambda x(1 - 3x)(1 + x^2)}{\eta(1 - x)^2} \quad (9)$$

This exhibits the crucial importance of the parameters  $\xi, x$ . Note that one can trade[12, 4, 19] the parameter  $\xi$  for  $x$  with advantage using the equation (8) since  $\xi$  is uniquely fixed by  $x$ . By a survey of the behaviour of the theory as a function of the complex parameter  $x$  we thus cover the behaviour of the three different solutions possible for each complex value of  $\xi$ . However the  $\xi$  parametrization is crucial to study the spontaneous CP violation case that is a principal focus of this paper.

### 2.1.1 Digression on Supergravity Potential

So far the vacuum of MSGUTs has been determined by extremizing the global susy potential by satisfying the F and D conditions[10, 11, 12]. This was justified since the supergravity corrections to  $F$  would be  $\sim M_X^4/M_P^2 \ll M_X^2$ , if  $M_X \sim 10^{16} GeV$  as is customary in Susy GUTs. However in view of the proposal[23] to raise the GUT scales to the vicinity of the Planck mass, by virtue of threshold corrections that modify the MSSM unification value, Supergravity corrections to the potential would surely be important. If moreover as proposed in [23] one could employ the MSSM singlets in the GUT fields themselves to break supersymmetry and to tune the superheavy contributions to the cosmological constant to zero, the minimization of the full supergravity potential becomes essential.

In the MSGUT and NMSGUT there are 5 MSSM singlet chiral fields namely  $p(1, 1, 1)_{210}, a(15, 1, 1)_{210}, \omega(15, 1, 3)_{210}, \sigma(\overline{10}, 1, 3)_{126}, \sigma(10, 1, 3)_{\overline{126}}$  (in terms of the  $SU(4) \times SU(2)_L \times SU(2)_R$  labels of the submultiplets they lie in). The MSSM singlet superpotential takes the form

$$\begin{aligned} W(a, p, \omega, \sigma, \bar{\sigma}) = & m(p^2 + 3a^2 + 6\omega^2) + 2\lambda(a^3 + 3p\omega^2 + 6a\omega^2) \\ & + (M + \eta(p + 3a - 6\omega))\sigma\bar{\sigma} + W_0 + W_{dyn} \end{aligned} \quad (10)$$

where we have included a constant term  $W_0$  and a dynamical superpotential[25] function of chiral  $SO(10)$  invariants formed from the the singlet fields to account for gaugino condensation and non trivial chiral moduli when the theory condenses in the UV. Such a superpotential was used to demonstrate the feasibility of (calculable) dynamical symmetry breaking in an asymptotically strong  $SU(2)$  toy model[25]. This superpotential is highly constrained by the holomorphy of the Supersymmetric gauge theory and can in principle be determined by arguments similar to those we used for the toy model[25]. We propose, in good time, to do the same for the MSGUT but for the moment we shall ignore it. No other renormalization of the superpotential is expected due to holomorphy[26].

We have deliberately not inserted any other  $SO(10)$  invariant non-renormalizable gravity induced terms in the superpotential since we expect that gravity itself may be induced by the UV condensation of the GUT. In the aftermath of the AdS-CFT conjecture the expectation of an effective induced gravity as a consequence of gauge

condensation has become almost passé [28]. However here no exotic extra dimensional context is involved but only the one context where supersymmetry can confidently be expected to be obeyed and a GUT that has passed a series of consistency tests and challenges to its accommodative abilities without falling into a disordered state of ad-hoc invocation of special mechanisms invoked to solve one-off problems. Predictability may well be retained since the induced gravity should have a potential of the  $N = 1$  supergravity form which is determined by the same unrenormalized (but dynamically corrected for gauge condensation) superpotential and possibly additional corrections to the kinetic terms by modification of the Kahler potential only[25]. If one was forced to include all possible or even some part of the possible gauge invariant but non-renormalizable superpotential operators we would lose all hope of a predictive or quantitative understanding that was not merely a multi-parameter re-formulation of our ignorance and hope. The scenario we advocate provides at least the skeleton of a rationale for the absence of all such arbitrary corrections due to the known power of holomorphy in protecting the superpotential.

The supergravity potential takes the form[30] ( $\kappa = (M_P^*)^{-1}$ ;  $M_P^* = M_P/\sqrt{8\pi} = 2 \times 10^{18} GeV$ )

$$\begin{aligned}
V &= -\mathcal{F}_i \mathcal{G}^i_j \mathcal{F}^{j*} - 3\kappa^{-4} e^{-\kappa^2 \mathcal{G}} + \sum \frac{g^2}{2} f_{ab}^{-1} D^a D^b \\
D^{\alpha a} &= \mathcal{G}^i (T^{\alpha a})_i^j \phi_j \\
\mathcal{G} &= -\sum_i \phi_i \phi^{i*} - \kappa^{-2} \log \kappa^6 |W(a, p, \omega, \sigma, \bar{\sigma})|^2 \\
\mathcal{F}_i &= \frac{1}{\kappa} e^{-\mathcal{G}\kappa^2/2} (\mathcal{G}^{-1})^j_i \mathcal{G}_j + \frac{1}{4} f_{ab}^* k (\mathcal{G}^{-1})^j_i \bar{\lambda}^a \bar{\lambda}^b
\end{aligned} \tag{11}$$

Where  $f_{ab}$  is the gauge kinetic function and the notation is standard[33]. Clearly the analysis of the full problem is quite daunting. Let us begin by writing down the potential in the simplified case where we take the kinetic terms to be canonical and ignore the dynamical superpotential and gaugino bilinears in  $\mathcal{F}$ . Then the supergravity F terms take the form ( $K = |p|^2 + 3|a|^2 + 6|\omega|^2 + \frac{|\sigma|^2 + |\bar{\sigma}|^2}{2}$ )

$$\begin{aligned}
\mathcal{F}_p &= (2mp + 6\lambda\omega^2 + \eta\sigma\bar{\sigma} + \kappa^2 p^* W) e^{-\kappa^2 K} \\
\mathcal{F}_a &= \sqrt{3}(2ma + 2\lambda(a^2 + 2\omega^2) + \eta\sigma\bar{\sigma} + \kappa^2 a^* W) e^{-\kappa^2 K} \\
\mathcal{F}_\omega &= \sqrt{6}(2m\omega + 2\omega\lambda(p + 2a) - \eta\sigma\bar{\sigma} + \kappa^2 \omega^* W) e^{-\kappa^2 K} \\
\mathcal{F}_\sigma &= \frac{1}{\sqrt{2}}((M + \eta(p + 3a - 6\omega)\bar{\sigma} + \kappa^2 \sigma^* W) e^{-\kappa^2 K} \\
\mathcal{F}_{\bar{\sigma}} &= \frac{1}{\sqrt{2}}((M + \eta(p + 3a - 6\omega)\sigma + \kappa^2 \bar{\sigma}^* W) e^{-\kappa^2 K}
\end{aligned} \tag{12}$$

$$V = e^{\kappa^2 K} \left( \sum_i |\mathcal{F}_i|^2 - 3\kappa^2 |W|^2 \right) + \frac{g^2}{2} \sum_a (D^a)^2 \tag{13}$$

Now the standard procedure[30, 23] can be followed : extremize  $\partial V/\partial(a, p, \omega, \sigma, \bar{\sigma}) = 0$ , impose  $V| = 0$  at the extremae and check that the mass spectrum is non tachyonic by appropriate conditions on the superpotential parameters. Since  $W$  is only cubic in the fields, the potential that must be minimized is effectively of degree 8. Although complicated this is definitely an improvement over a quintic[23] superpotential (with many unwanted relatives waiting in the the undergrowth for the door to be opened!). The multiplicity of singlets effectively raises the degree without introducing new parameters.

Assuming the D terms continue to vanish separately (only  $|\sigma| = |\bar{\sigma}|$  is needed) and the vacuum energy at the minimum is tuned to zero, the equations which must be solved in the case of a cubic superpotential are ( $\partial_i \equiv \partial/\partial\phi^i$ )

$$\partial_i \partial_j W \partial_j^* W^* + \kappa^2 \partial_i \partial_j W \phi^j W^* + \kappa^2 \partial_i W (W^{(3)} + \kappa^2 K W)^* + \kappa^4 \phi^* |W|^2 = 0 \quad (14)$$

Here  $W^{(3)}$  is the part of the superpotential cubic in the fields. While a detailed analysis must be deferred[35] it is clear that this potential has 5 free parameters i.e  $W_0, \lambda, \eta, m, M$  and these could well be sufficient to obtain a solution of the desired type i.e with vevs  $\leq \kappa^{-1}$  and  $m_{\frac{3}{2}} \ll M_X$  since the raising of gauge scale and the baryon decay mediating triplet will be found(see below) to be practically automatic. We now continue with the analysis of the theory with the global susy potential with a particular view to show that threshold corrections tend to raise the unification scale.

### 2.1.2 Characteristics of the SSB solutions

A knowledge of the 3 solutions  $x_i(\xi)(i = 1, 2, 3 \mid C(x_i(\xi), \xi) = 0)$  of the cubic equation(8) is necessary when considering issues such as the viability of spontaneous(GUT scale) CP violation in this theory.

In Fig. 1,2 we exhibit plots of  $x_i(\xi)$  for *real*  $\xi$ . For  $\xi < -27.917$  all three solutions are real. Moreover it is clear that for  $|x| \gg 1$ ,  $x \simeq -\xi/8$  specifies one real branch for real  $\xi$  as visible in Fig. 1. On the other two (complex conjugate) branches the real and imaginary parts of  $x(\xi)$  are bounded above and below( real part  $\in (.8, 1.0)$  and imaginary part magnitude  $\in [0, 1.1)$ ).

Using the above vevs and the methods of [3] we calculated the complete gauge and chiral multiplet GUT scale spectra *and* couplings for the 52 different MSSM multiplet sets falling into 26 different MSSM multiplet types (prompting a natural alphabetization of their naming convention[3]) of which 18 are unmixed while the other 8 types occur in multiple copies which mix. The (full details of these) spectra may be found in[3, 5] and equivalent results(with slightly differing conventions ) are presented in[4]. A related calculation with very different conventions has been reported in [6]. The initially controversial relation between the overlapping parts of these papers was discussed and resolved in [20].






Figure 1: Solutions of eqn.(8) which governs GUT ssb: Plot of  $Re[x_i(\xi)]$  vs  $\xi$  for  $i = 1, 2, 3$ . The vertical straight line segments are “reconnection artifacts” induced by a switch over between real and complex solutions and vice versa.

Among the mass matrices is the all important  $4 \times 4$  Higgs doublet mass matrix [3, 5]  $\mathcal{H}$  which can be diagonalized by a bi-unitary transformation[12, 4, 5]: from the 4 pairs of Higgs doublets  $h^{(i)}, \bar{h}^{(i)}$  arising from the SO(10) fields to a new set  $H^{(i)}, \bar{H}^{(i)}$  of fields in terms of which the doublet mass terms are diagonal.

$$\begin{aligned} \bar{U}^T \mathcal{H} U &= \text{Diag}(m_H^{(1)}, m_H^{(2)}, \dots) \\ h^{(i)} &= U_{ij} H^{(j)} \quad ; \quad \bar{h}^{(i)} = \bar{U}_{ij} \bar{H}^{(j)} \end{aligned} \quad (15)$$

To keep one pair of these doublets light one tunes  $M_H$  so that  $\text{Det} \mathcal{H} = 0$ . In the effective theory at low energies the GUT Higgs doublets  $h^{(i)}, \bar{h}^{(i)}$  are present in the massless doublets  $H^{(1)}, \bar{H}^{(1)}$  in a proportion determined by the first columns of the matrices  $U, \bar{U}$  :

$$\begin{aligned} E \ll M_X \quad : \quad h^{(i)} &\rightarrow \alpha_i H^{(1)} \quad ; \quad \alpha_i = U_{i1} \\ \bar{h}^{(i)} &\rightarrow \bar{\alpha}_i \bar{H}^{(1)} \quad ; \quad \bar{\alpha}_i = \bar{U}_{i1} \end{aligned} \quad (16)$$

The all important normalized 4-tuples  $\alpha, \bar{\alpha}$  can be easily determined[12, 4, 14, 19, 15] by solving the zero mode conditions:  $\mathcal{H}\alpha = 0$  ;  $\bar{\alpha}^T \mathcal{H} = 0$ .




Figure 2: Solutions of eqn.(8) which governs GUT ssb: Plot of  $Im[x_i(\xi)]$  vs  $\xi$  for  $i = 1, 2, 3$ . The vertical straight line segments are again ‘reconnection artifacts’.

## 2.2 Additional terms introduced by the 120

The introduction of the **120**-plet Higgs representation leads to new couplings in the superpotential . The additional terms are

$$\begin{aligned}
 W_{NMSG} = & \frac{m_o}{2(3!)} O_{ijk} O_{ijk} + \frac{k}{3!} O_{ijk} H_m \Phi_{mijk} + \frac{\rho}{4!} O_{ijk} O_{mnk} \Phi_{ijmn} \\
 & + \frac{1}{2(3!)} O_{ijk} \Phi_{klmn} (\zeta \Sigma_{lmnij} + \bar{\zeta} \bar{\Sigma}_{lmnij}) + \frac{1}{5!} g_{AB} \Psi_A^T C_2^{(5)} \gamma_{i_1} \gamma_{i_2} \gamma_{i_3} \Psi_B O_{i_1 i_2 i_3}
 \end{aligned}$$

The Yukawa coupling  $g_{AB}$  is a complex antisymmetric  $3 \times 3$  matrix. The  $SU(4) \times SU(2)_L \times SU(2)_R$  (Pati-Salam) decomposition of the **120**plet is as follows :

$$\begin{aligned}
 O_{ijk}(120) = & O_{\mu\nu}^{(s)}(10, 1, 1) + \bar{O}_{(s)}^{\mu\nu}(\bar{10}, 1, 1) + O_{\nu\alpha\dot{\alpha}}^{\mu}(15, 2, 2) \\
 & + O_{\mu\nu\dot{\alpha}\dot{\beta}}^{(a)}(6, 1, 3) + O_{\mu\nu\alpha\beta}^{(a)}(6, 3, 1) + O_{\alpha\dot{\alpha}}(1, 2, 2)
 \end{aligned} \tag{17}$$

The arbitrary phase of the **120** reduces the effective number of the extra couplings ( $m_o, \rho, \zeta, \bar{\zeta}, k$  ( $5_c - 1_r = 9_r$ ) and  $g_{AB}$  ( $3_c = 6_r$ )) so they amount to 15 additional parameters. Thus the relative advantage[12, 29] with respect to  $SU(5)$  theories using

additional fields or higher dimensional operators to correct the fermion mass relations of the simplest  $SU(5)$  model seems weakened. *However this is too hasty a conclusion.*

In fact the old MSGUT fails to fit the fermion mass data due to difficulties with the overall neutrino mass scale[14, 15]. An alternative scenario within the NMSGUT which successfully evades the problems of the old MSGUT was proposed and elaborated in [15, 16, 17]. In this scenario the Yukawa couplings ( $f_{AB}$ ) of the  $\overline{\mathbf{126}}$  are much smaller than those of the  $\mathbf{10}, \mathbf{120}$ . This boosts the value of the Type I seesaw masses (which were in any case dominant over the Type II seesaw masses but still too small) so that they are comfortably generically viable. We demonstrated the feasibility of this fitting ansatz in CP preserving 2 and 3 generation models using a perturbation around the dominant 2-3 sector of the fermion mass matrices. Furthermore, using a “downhill simplex ” non linear fitting procedure the authors of [18] were able to show that even with CP preserving (i.e real) Yukawa couplings the CKM CP violation as well as all the other known fermion mass and mixing data could be accurately fit with Yukawa couplings which followed the ‘ $\overline{\mathbf{126}}$  suppressed’ dominance pattern. Although the Yukawa coupling of the  $\mathbf{10}$  plet Higgs (weakly) dominates that of the  $\overline{\mathbf{126}}$  in the solution of [18], the same is not true of the  $\mathbf{120}$ . This is rather different from the situation in the perturbative fit obtained by us[17] where the couplings of the  $\overline{\mathbf{126}}$  are very small. Since these two solutions are only early attempts to view the vast parameter space of the fit we shall not belabour the difference.

The authors of [18] fixed 3 of the Yukawa parameters using an ad hoc  $Z_2$  symmetry. Thus these 3 parameters remain “in reserve” and can be called to fit the values of the as yet unknown Leptonic CP phases and Neutrino mass scale besides the actual value of  $\theta_{13}^P$  (as opposed to a limit). The indications are thus that even a NMSGUT with only real Yukawas i.e with a total of 24(= 12 fermion Yukawas + 12 AM Yukawas) real parameters (further reduced to 23 by the fine tuning condition) will still be able to fit all the fermion data ( 12 masses + CKM phase + 3 CKM angles + 3 PMNS angles + 3 PMNS phases = 22 parameters) ! Such a theory will have less parameters than even the (unsuccessful) old MSGUT ! Moreover this fitting problem which will have 22 data values fit by at most 23 parameters (once the missing neutrino data is available and counting phases as “full” parameters !) holds out the enticing prospect of predictions regarding ‘smoking gun ’ baryon violation signals of grand unification. Even before such data is available however, it would clearly be preferable to analyze the simpler case of spontaneous CP violation first. This shall be a guiding theme in our discussions of the NMSGUT parameter space.

It is clear that that the  $\mathbf{120}$  does not contain any SM singlets and hence the analysis of the GUT scale symmetry breaking in MSGUT carries over unchanged to the NMSGUT. In particular there is still only one complex parameter( $x$ ) whose variation directly affects the vevs and thus the masses in the theory. The additional kinetic terms are given by covariantizing in the standard way the global  $SO(10)$  invariant D-terms  $[\frac{1}{(3)!}O_{ijk}^*O_{ijk}]_D$

### 3 AM Chiral masses via PS

As in the case of the MSGUT [3, 5] we open up the maze of NMSGUT interactions by decomposing  $SO(10)$  invariants in the superpotential first into Pati-Salam invariants and then, after substituting the GUT scale vevs in PS notation we obtain the superpotential in the MSSM vacuum in terms of MSSM invariants. The results for the old MSGUT case were already given in [3, 5] thus we list only the effect of the additional terms in the superpotential. The PS form of  $W_{NMSG}$  is (we have inserted line numbers for easy reference) :

$$\frac{k}{3!}H_i O_{jkl} \Phi_{ijkl} = k[\frac{1}{\sqrt{2}}i(\tilde{H}_{(a)}^{\mu\nu} O_{\mu\lambda}^{(s)} \Phi_{\nu}^{\lambda} + H_{\mu\nu}^{(a)} O_{(s)}^{\mu\lambda} \Phi_{\lambda}^{\nu}) \quad (18)$$

$$+ \frac{1}{\sqrt{2}}(\tilde{H}_{(a)}^{\mu\nu} O_{\nu}^{\lambda\alpha\dot{\alpha}} \Phi_{\mu\lambda\alpha\dot{\alpha}}^{(s)} - H_{\mu\nu}^{(a)} O_{\lambda}^{\nu\alpha\dot{\alpha}} \Phi_{\alpha\dot{\alpha}}^{\mu\lambda(s)}) \quad (19)$$

$$+ H_{\mu\nu}^{(a)} (\vec{O}_{(a)(R)}^{\nu\lambda} \cdot \vec{\Phi}_{\lambda(R)}^{\mu} + \vec{O}_{(a)(L)}^{\nu\lambda} \cdot \vec{\Phi}_{\lambda(L)}^{\mu}) \quad (20)$$

$$- \frac{1}{2} \tilde{H}_{(a)}^{\mu\nu} O_{\mu\nu\alpha\dot{\alpha}}^{\alpha\dot{\alpha}} \Phi_{\mu\nu\alpha\dot{\alpha}}^{(a)} \quad (21)$$

$$+ \frac{1}{2} H^{\alpha\dot{\alpha}} (O_{\mu\nu}^{(s)} \Phi_{\alpha\dot{\alpha}}^{\mu\nu(s)} + O_{(s)}^{\mu\nu} \Phi_{\mu\nu\alpha\dot{\alpha}}^{(s)}) \quad (22)$$

$$- \frac{1}{\sqrt{2}} H^{\alpha\dot{\alpha}} (O_{\nu\alpha}^{\mu\dot{\beta}} \Phi_{\mu\dot{\alpha}\dot{\beta}}^{\nu} + O_{\nu\dot{\alpha}}^{\mu\beta} \Phi_{\mu\alpha\beta}^{\nu}) \quad (23)$$

$$- \frac{1}{2\sqrt{2}} H^{\alpha\dot{\alpha}} (\tilde{O}_{(a)\dot{\alpha}\dot{\beta}}^{\mu\nu} \Phi_{\mu\nu\alpha}^{(a)\dot{\beta}} - \tilde{O}_{(a)\alpha\beta}^{\mu\nu} \Phi_{\mu\nu\dot{\alpha}}^{(a)\beta}) \quad (24)$$

$$+ H^{\alpha\dot{\alpha}} O_{\alpha\dot{\alpha}} \Phi] \quad (25)$$

$$\frac{m_0}{2(3!)} O_{ijk} O_{ijk} = \frac{m_0}{12} [6 O_{\mu\nu}^{(s)} O_{(s)}^{\mu\nu} + 6 O_{\sigma}^{\lambda\alpha\dot{\alpha}} O_{\lambda\alpha\dot{\alpha}}^{\sigma} \quad (26)$$

$$+ 3(\vec{O}_{(a)(R)}^{\mu\nu} \cdot \vec{O}_{\mu\nu(R)}^{(a)} + \vec{O}_{(a)(L)}^{\mu\nu} \cdot \vec{O}_{\mu\nu(L)}^{(a)}) \quad (27)$$

$$- 6 O^{\alpha\dot{\alpha}} O_{\alpha\dot{\alpha}}] \quad (28)$$

$$\frac{\rho}{4!} O_{ijm} O_{klm} \Phi_{ijkl} = \frac{\rho}{4!} [8i O_{\mu\lambda}^{(s)} O_{(s)}^{\nu\lambda} \Phi_{\nu}^{\mu} + 8i O_{\sigma}^{\mu\alpha\dot{\alpha}} O_{\nu\alpha\dot{\alpha}}^{\sigma} \Phi_{\mu}^{\nu} \quad (29)$$

$$- 4\sqrt{2} (O_{\mu\lambda}^{(s)} \vec{O}_{(a)(R)}^{\mu\nu} \cdot \vec{\Phi}_{\nu(R)}^{\lambda} + O_{\mu\lambda}^{(s)} \vec{O}_{(a)(L)}^{\mu\nu} \cdot \vec{\Phi}_{\nu(L)}^{\lambda}) \quad (30)$$

$$- O_{(s)}^{\mu\lambda} \vec{O}_{\mu\nu(R)}^{(a)} \cdot \vec{\Phi}_{\lambda(R)}^{\nu} - O_{(s)}^{\mu\lambda} \vec{O}_{\mu\nu(L)}^{(a)} \cdot \vec{\Phi}_{\lambda(L)}^{\nu}) \quad (31)$$

$$+ 4\sqrt{2} (O^{\alpha\dot{\alpha}} O_{\nu\alpha}^{\mu\dot{\beta}} \Phi_{\mu\dot{\alpha}\dot{\beta}}^{\nu} - O^{\alpha\dot{\alpha}} O_{\nu\dot{\alpha}}^{\mu\beta} \Phi_{\mu\alpha\beta}^{\nu}) \quad (32)$$

$$+ 8 (O_{\nu}^{\mu\alpha\dot{\alpha}} O_{\mu\lambda}^{(s)} \Phi_{\alpha\dot{\alpha}}^{\nu\lambda(s)} - O_{\nu}^{\mu\alpha\dot{\alpha}} O_{(s)}^{\nu\lambda} \Phi_{\mu\lambda\alpha\dot{\alpha}}^{(s)}) \quad (33)$$

$$- 2(\tilde{O}_{(a)\dot{\alpha}\dot{\beta}}^{\mu\nu} O_{\nu\alpha}^{\lambda\dot{\beta}} \Phi_{\mu\lambda(s)}^{\alpha\dot{\alpha}} + \tilde{O}_{(a)\alpha\beta}^{\mu\nu} O_{\nu\dot{\alpha}}^{\lambda\beta} \Phi_{\mu\lambda(s)}^{\alpha\dot{\alpha}}) \quad (34)$$

$$-O_{\mu\nu\dot{\alpha}\dot{\beta}}^{(a)} O_{\lambda\alpha}^{\nu\dot{\beta}} \Phi_{(s)}^{\mu\lambda\alpha\dot{\alpha}} - O_{\mu\nu\alpha\beta}^{(a)} O_{\lambda\dot{\alpha}}^{\nu\beta} \Phi_{(s)}^{\mu\lambda\alpha\dot{\alpha}} \quad (35)$$

$$-2(\vec{O}_{(a)(L)}^{\mu\nu} \cdot \vec{O}_{\mu\nu(L)}^{(a)} - \vec{O}_{(a)(R)}^{\mu\nu} \cdot \vec{O}_{\mu\nu(R)}^{(a)}) \Phi \quad (36)$$

$$-4\sqrt{2}(O_{\lambda}^{\mu\alpha\dot{\alpha}} O_{(a)\dot{\alpha}\dot{\beta}}^{\nu\lambda} \Phi_{\mu\nu\alpha}^{(a)\dot{\beta}} - O_{\lambda}^{\mu\alpha\dot{\alpha}} O_{(a)\alpha\beta}^{\nu\lambda} \Phi_{\mu\nu\dot{\alpha}}^{(a)\beta}) \quad (37)$$

$$-2\sqrt{2}(O^{\alpha\dot{\alpha}} \tilde{O}_{(a)\dot{\alpha}\dot{\beta}}^{\mu\nu} \Phi_{\mu\nu\alpha}^{(a)\dot{\beta}} + O^{\alpha\dot{\alpha}} \tilde{O}_{(a)\alpha\beta}^{\mu\nu} \Phi_{\mu\nu\dot{\alpha}}^{(a)\beta}) \quad (38)$$

$$+4\sqrt{2}(O_{\lambda}^{\nu\alpha\dot{\alpha}} O_{\mu\alpha}^{\lambda\dot{\beta}} \Phi_{\nu\dot{\alpha}\dot{\beta}}^{\mu} + O_{\lambda}^{\nu\alpha\dot{\alpha}} O_{\mu\dot{\alpha}}^{\lambda\beta} \Phi_{\nu\alpha\beta}^{\mu}) \quad (39)$$

$$-2(O_{\nu\alpha}^{\lambda\dot{\beta}} \tilde{O}_{\dot{\alpha}\dot{\beta}}^{\mu\nu(a)} \Phi_{\mu\lambda(s)}^{\alpha\dot{\alpha}} + O_{\nu\dot{\alpha}}^{\lambda\beta} \tilde{O}_{\alpha\beta}^{\mu\nu(a)} \Phi_{\mu\lambda(s)}^{\alpha\dot{\alpha}}) \quad (40)$$

$$-O_{\lambda\dot{\alpha}}^{\nu\beta} O_{\mu\nu\alpha\beta}^{(a)} \Phi_{(s)}^{\mu\lambda\alpha\dot{\alpha}} - O_{\lambda\alpha}^{\nu\dot{\beta}} O_{\mu\nu\dot{\alpha}\dot{\beta}}^{(a)} \Phi_{(s)}^{\mu\lambda\alpha\dot{\alpha}} \quad (41)$$

$$-4\sqrt{2}(\vec{O}_{\mu\nu(R)}^{(a)} \cdot (\vec{O}_{\nu\lambda}^{\mu} \times \vec{\Phi}_{\lambda(R)}^{\mu})) \quad (42)$$

$$+ \vec{O}_{\mu\nu(L)}^{(a)} \cdot (\vec{O}_{\nu\lambda}^{\mu} \times \vec{\Phi}_{\lambda(L)}^{\mu})) \quad (43)$$

$$\frac{\zeta}{2(3!)} O_{ijk} \Sigma_{ijlmn} \Phi_{klmn} = \frac{\zeta}{2(3!)} [-6\sqrt{2}i(O_{\mu\lambda}^{(s)} \tilde{\Sigma}_{(a)}^{\mu\nu} \Phi_{\nu}^{\lambda} - O_{(s)}^{\mu\lambda} \Sigma_{\mu\nu}^{(a)} \Phi_{\lambda}^{\nu}) \quad (44)$$

$$-12i(O_{\mu\lambda}^{(s)} \Sigma_{\nu}^{\mu\alpha\dot{\alpha}} \Phi_{\alpha\dot{\alpha}}^{\nu\lambda(s)} + O_{(s)}^{\nu\lambda} \Sigma_{\nu}^{\mu\alpha\dot{\alpha}} \Phi_{\mu\lambda\alpha\dot{\alpha}}^{(s)}) \quad (45)$$

$$-12\sqrt{2}(O_{\mu\lambda}^{(s)} \tilde{\Sigma}_{(s)(R)}^{\nu\lambda} \cdot \vec{\Phi}_{\nu(R)}^{\mu} - O_{(s)}^{\nu\lambda} \tilde{\Sigma}_{\mu\lambda(L)}^{\nu\lambda} \cdot \vec{\Phi}_{\nu(L)}^{\mu}) \quad (46)$$

$$+6\sqrt{2}i(O_{(s)}^{\mu\lambda} \Sigma_{\lambda\alpha\dot{\alpha}}^{\nu} \Phi_{\mu\nu(a)}^{\alpha\dot{\alpha}} - O_{\mu\lambda}^{(s)} \Sigma_{\nu\alpha\dot{\alpha}}^{\lambda} \tilde{\Phi}_{(a)}^{\mu\nu\alpha\dot{\alpha}}) \quad (47)$$

$$-6\sqrt{2}(O_{\nu}^{\lambda\alpha\dot{\alpha}} \tilde{\Sigma}_{(a)}^{\mu\nu} \Phi_{\mu\lambda\alpha\dot{\alpha}}^{(s)} + O_{\lambda}^{\nu\alpha\dot{\alpha}} \Sigma_{\mu\nu}^{(a)} \Phi_{\alpha\dot{\alpha}}^{\mu\lambda(s)}) \quad (48)$$

$$-12\sqrt{2}i(O_{\lambda}^{\mu\alpha\dot{\alpha}} \Sigma_{\nu\dot{\alpha}}^{\lambda\beta} \Phi_{\mu\alpha\beta}^{\nu} + O_{\lambda}^{\mu\alpha\dot{\alpha}} \Sigma_{\mu\alpha}^{\nu\beta} \Phi_{\nu\dot{\alpha}\dot{\beta}}^{\lambda}) \quad (49)$$

$$+6\sqrt{2}(O_{\nu\dot{\alpha}}^{\lambda\beta} \Sigma_{\mu\lambda\alpha\beta}^{(s)} \tilde{\Phi}_{(a)}^{\mu\nu\alpha\dot{\alpha}} + O_{\lambda\alpha}^{\nu\dot{\beta}} \Sigma_{\dot{\alpha}\dot{\beta}}^{\mu\lambda(s)} \Phi_{\mu\nu(a)}^{\alpha\dot{\alpha}}) \quad (50)$$

$$+12iO_{\nu}^{\mu\alpha\dot{\alpha}} \Sigma_{\mu\alpha\dot{\alpha}}^{\nu} \Phi \quad (51)$$

$$+12i(\vec{O}_{(a)(L)}^{\mu\nu} \cdot \vec{\Sigma}_{\mu\lambda(L)}^{(s)} \Phi_{\nu}^{\lambda} + \vec{O}_{\mu\nu(R)}^{(a)} \cdot \vec{\Sigma}_{(s)(R)}^{\mu\lambda} \Phi_{\lambda}^{\nu}) \quad (52)$$

$$-6i(\tilde{O}_{(a)\dot{\alpha}\dot{\beta}}^{\mu\nu} \Sigma_{\nu\alpha}^{\lambda\dot{\beta}} \Phi_{\mu\lambda(s)}^{\alpha\dot{\alpha}} - \tilde{O}_{(a)\alpha\beta}^{\mu\nu} \Sigma_{\nu\dot{\alpha}}^{\lambda\beta} \Phi_{\mu\lambda(s)}^{\alpha\dot{\alpha}}) \quad (53)$$

$$-O_{\mu\nu\dot{\alpha}\dot{\beta}}^{(a)} \Sigma_{\lambda\alpha}^{\nu\dot{\beta}} \Phi_{(s)}^{\mu\lambda\alpha\dot{\alpha}} + O_{\mu\nu\alpha\beta}^{(a)} \Sigma_{\lambda\dot{\alpha}}^{\nu\beta} \Phi_{(s)}^{\mu\lambda\alpha\dot{\alpha}} \quad (54)$$

$$-12(\vec{O}_{\mu\nu(L)}^{(a)} \vec{\Phi}_{\lambda(L)}^{\mu} - \vec{O}_{\mu\nu(R)}^{(a)} \vec{\Phi}_{\lambda(R)}^{\mu}) \Sigma_{(a)}^{\nu\lambda} \quad (55)$$

$$+6(O_{\dot{\alpha}}^{\beta} \Sigma_{\mu\nu\alpha\beta}^{(s)} \Phi_{(s)}^{\mu\nu\alpha\dot{\alpha}} - O_{\alpha}^{\dot{\beta}} \Sigma_{\dot{\alpha}\dot{\beta}}^{\mu\nu(s)} \Phi_{\mu\nu(s)}^{\alpha\dot{\alpha}}) \quad (56)$$

$$-6\sqrt{2}i(O_{\alpha}^{\dot{\beta}} \Sigma_{\nu}^{\mu\alpha\dot{\alpha}} \Phi_{\mu\dot{\alpha}\dot{\beta}}^{\nu} + O_{\dot{\alpha}}^{\beta} \Sigma_{\nu}^{\mu\alpha\dot{\alpha}} \Phi_{\mu\alpha\beta}^{\nu}) \quad (57)$$

$$+12(O_{\nu}^{\mu\alpha\dot{\alpha}} \Sigma_{\lambda\alpha\dot{\alpha}}^{\nu} \Phi_{\mu}^{\lambda} - O_{\nu}^{\mu\alpha\dot{\alpha}} \Sigma_{\mu\alpha\dot{\alpha}}^{\lambda} \Phi_{\lambda}^{\nu}) \quad (58)$$

$$+12(O_{\nu}^{\mu\alpha\dot{\alpha}} \Sigma_{\mu\lambda\alpha\beta}^{(s)} \Phi_{(s)\dot{\alpha}}^{\nu\lambda\beta} - O_{\nu}^{\mu\alpha\dot{\alpha}} \Sigma_{(s)\dot{\alpha}\dot{\beta}}^{\nu\lambda} \Phi_{\mu\lambda\alpha}^{(s)\dot{\beta}}) \quad (59)$$

$$-12(O_{\lambda}^{\mu\alpha\dot{\alpha}} \Sigma_{\mu\nu}^{(a)} \Phi_{(a)\alpha\dot{\alpha}}^{\nu\lambda}) \quad (60)$$

$$+6i(\tilde{O}_{(a)\dot{\alpha}\dot{\beta}}^{\mu\nu} \Sigma_{\nu}^{\lambda\alpha\dot{\alpha}} \Phi_{\mu\lambda\alpha}^{(s)\dot{\beta}} + \tilde{O}_{(a)\alpha\beta}^{\mu\nu} \Sigma_{\nu}^{\lambda\alpha\dot{\alpha}} \Phi_{\mu\lambda\dot{\alpha}}^{(s)\beta}) \quad (61)$$

$$+O_{\mu\nu\dot{\alpha}\dot{\beta}}^{(a)}\Sigma_{\lambda}^{\nu\alpha\dot{\alpha}}\Phi_{(s)\alpha}^{\mu\lambda\dot{\beta}}+O_{\mu\nu\alpha\beta}^{(a)}\Sigma_{\lambda}^{\nu\alpha\dot{\alpha}}\Phi_{(s)\dot{\alpha}}^{\mu\lambda\beta}) \quad (62)$$

$$+12\sqrt{2}[\vec{O}_{(a)(L)}^{\mu\nu}\cdot(\vec{\Phi}_{\nu(L)}^{\lambda}\times\vec{\Sigma}_{\mu\lambda(L)}^{(s)})] \quad (63)$$

$$-\vec{O}_{\mu\nu(R)}^{(a)}\cdot(\vec{\Phi}_{\lambda(R)}^{\nu}\times\vec{\Sigma}_{(R)}^{\mu\lambda(s)})] \quad (64)$$

$$+6\sqrt{2}i(O_{\mu\nu\dot{\alpha}\dot{\beta}}^{(a)}\Sigma_{\lambda\dot{\alpha}}^{\mu\dot{\beta}}\Phi_{(a)}^{\nu\lambda\alpha\dot{\alpha}}+O_{\mu\nu\alpha\beta}^{(a)}\Sigma_{\lambda\dot{\alpha}}^{\mu\beta}\Phi_{(a)}^{\nu\lambda\alpha\dot{\alpha}})] \quad (65)$$

$$\frac{\bar{\zeta}}{2(3!)}O_{ijk}\bar{\Sigma}_{ijlmn}\Phi_{klmn} = \frac{\bar{\zeta}}{2(3!)}[6\sqrt{2}i(O_{\mu\lambda}^{(s)}\bar{\Sigma}_{\nu}^{\mu\lambda}\Phi_{\nu}^{\lambda}-O_{(s)}^{\mu\lambda}\bar{\Sigma}_{\mu\nu}^{(a)}\Phi_{\lambda}^{\nu}) \quad (66)$$

$$-12i(O_{\mu\lambda}^{(s)}\bar{\Sigma}_{\nu}^{\mu\alpha\dot{\alpha}}\Phi_{\alpha\dot{\alpha}}^{\nu\lambda(s)}+O_{(s)}^{\nu\lambda}\bar{\Sigma}_{\nu}^{\mu\alpha\dot{\alpha}}\Phi_{\mu\lambda\alpha\dot{\alpha}}^{(s)}) \quad (67)$$

$$-12\sqrt{2}(O_{\mu\lambda}^{(s)}\bar{\Sigma}_{\nu(L)}^{\mu\lambda(s)}\cdot\vec{\Phi}_{\nu(L)}^{\mu}-O_{(s)}^{\nu\lambda}\bar{\Sigma}_{\mu\lambda(R)}^{\mu\lambda(s)}\cdot\vec{\Phi}_{\nu(R)}^{\mu}) \quad (68)$$

$$-6\sqrt{2}i(O_{(s)}^{\mu\lambda}\bar{\Sigma}_{\lambda\alpha\dot{\alpha}}^{\nu}\Phi_{\mu\nu}^{(a)\alpha\dot{\alpha}}-O_{\mu\lambda}^{(s)}\bar{\Sigma}_{\nu\alpha\dot{\alpha}}^{\lambda}\Phi_{(a)}^{\mu\nu\alpha\dot{\alpha}}) \quad (69)$$

$$+6\sqrt{2}(O_{\nu}^{\lambda\alpha\dot{\alpha}}\bar{\Sigma}_{(a)}^{\mu\nu}\Phi_{\mu\lambda\alpha\dot{\alpha}}^{(s)}+O_{\lambda}^{\nu\alpha\dot{\alpha}}\bar{\Sigma}_{\mu\nu}^{(a)}\Phi_{\mu\lambda\alpha\dot{\alpha}}^{(s)}) \quad (70)$$

$$-12\sqrt{2}i(O_{\lambda}^{\mu\alpha\dot{\alpha}}\bar{\Sigma}_{\nu\alpha}^{\lambda\dot{\beta}}\Phi_{\mu\dot{\alpha}\dot{\beta}}^{\nu}+O_{\lambda}^{\mu\alpha\dot{\alpha}}\bar{\Sigma}_{\mu\dot{\alpha}}^{\nu\beta}\Phi_{\nu\alpha\beta}^{\lambda}) \quad (71)$$

$$-6\sqrt{2}(O_{\nu\alpha}^{\lambda\dot{\beta}}\bar{\Sigma}_{\mu\dot{\alpha}\dot{\beta}}^{(s)}\Phi_{(a)}^{\mu\nu\alpha\dot{\alpha}}+O_{\lambda\dot{\alpha}}^{\nu\beta}\bar{\Sigma}_{\alpha\beta}^{\mu\lambda(s)}\Phi_{\mu\nu}^{(a)\alpha\dot{\alpha}}) \quad (72)$$

$$-12iO_{\nu}^{\mu\alpha\dot{\alpha}}\bar{\Sigma}_{\mu\alpha\dot{\alpha}}^{\nu}\Phi \quad (73)$$

$$+12i(\vec{O}_{(R)}^{\mu\nu(a)}\cdot\vec{\Sigma}_{\mu\lambda(R)}^{(s)}\Phi_{\nu}^{\lambda}+\vec{O}_{\mu\nu(L)}^{(a)}\cdot\vec{\Sigma}_{(L)}^{\mu\lambda(s)}\Phi_{\lambda}^{\nu}) \quad (74)$$

$$+6i(\tilde{O}^{\mu\nu(a)}_{\dot{\alpha}\dot{\beta}}\bar{\Sigma}_{\nu\alpha}^{\lambda\dot{\beta}}\Phi_{\mu\lambda(a)}^{\alpha\dot{\alpha}}-\tilde{O}^{\mu\nu(a)}_{\alpha\beta}\bar{\Sigma}_{\nu\dot{\alpha}}^{\lambda\dot{\beta}}\Phi_{\mu\lambda(a)}^{\alpha\dot{\alpha}}) \quad (75)$$

$$-O_{\mu\nu\dot{\alpha}\dot{\beta}}^{(a)}\bar{\Sigma}_{\lambda\alpha}^{\nu\dot{\beta}}\Phi_{(a)}^{\mu\lambda\alpha\dot{\alpha}}+O_{\mu\nu\alpha\beta}^{(a)}\bar{\Sigma}_{\lambda\dot{\alpha}}^{\nu\beta}\Phi_{(a)}^{\mu\lambda\alpha\dot{\alpha}}) \quad (76)$$

$$-12(\vec{O}_{\mu\nu(L)}^{(a)}\cdot\vec{\Phi}_{\lambda(L)}^{\mu}-\vec{O}_{\mu\nu(R)}^{(a)}\cdot\vec{\Phi}_{\lambda(R)}^{\mu})\bar{\Sigma}_{(a)}^{\nu\lambda} \quad (77)$$

$$-6(O_{\alpha}^{\dot{\beta}}\bar{\Sigma}_{\mu\nu\dot{\alpha}\dot{\beta}}^{(s)}\Phi_{(s)}^{\mu\nu\alpha\dot{\alpha}}-O_{\dot{\alpha}}^{\beta}\bar{\Sigma}_{\alpha\beta}^{\mu\nu(s)}\Phi_{\mu\nu(s)}^{\alpha\dot{\alpha}}) \quad (78)$$

$$-6\sqrt{2}i(O_{\alpha}^{\dot{\beta}}\bar{\Sigma}_{\nu}^{\mu\alpha\dot{\alpha}}\Phi_{\mu\dot{\alpha}\dot{\beta}}^{\nu}+O_{\dot{\alpha}}^{\beta}\bar{\Sigma}_{\nu}^{\mu\alpha\dot{\alpha}}\Phi_{\mu\alpha\beta}^{\nu}) \quad (79)$$

$$+12(O_{\nu}^{\mu\alpha\dot{\alpha}}\bar{\Sigma}_{\lambda\alpha\dot{\alpha}}^{\nu}\Phi_{\mu}^{\lambda}-O_{\nu}^{\mu\alpha\dot{\alpha}}\bar{\Sigma}_{\mu\alpha\dot{\alpha}}^{\lambda}\Phi_{\lambda}^{\nu}) \quad (80)$$

$$+12(O_{\nu}^{\mu\alpha\dot{\alpha}}\bar{\Sigma}_{\mu\lambda\dot{\alpha}\dot{\beta}}^{(s)}\Phi_{\alpha}^{\nu\lambda(s)\dot{\beta}}-O_{\nu}^{\mu\alpha\dot{\alpha}}\bar{\Sigma}_{\alpha\beta}^{\nu\lambda(s)}\Phi_{\mu\lambda(s)\dot{\alpha}}^{\beta}) \quad (81)$$

$$-12(O_{\lambda}^{\mu\alpha\dot{\alpha}}\bar{\Sigma}_{\mu\nu}^{(a)}\Phi_{(a)\alpha\dot{\alpha}}^{\nu\lambda}) \quad (82)$$

$$+6i(\tilde{O}_{(a)\dot{\alpha}\dot{\beta}}^{\mu\nu}\bar{\Sigma}_{\nu}^{\lambda\alpha\dot{\alpha}}\Phi_{\mu\lambda\alpha}^{(s)\dot{\beta}}+\tilde{O}_{(a)\alpha\beta}^{\mu\nu}\bar{\Sigma}_{\nu}^{\lambda\alpha\dot{\alpha}}\Phi_{\mu\lambda\dot{\alpha}}^{(s)\beta}) \quad (83)$$

$$+O_{\mu\nu\dot{\alpha}\dot{\beta}}^{(a)}\bar{\Sigma}_{\lambda}^{\nu\alpha\dot{\alpha}}\Phi_{\alpha}^{\mu\lambda(s)\dot{\beta}}+O_{\mu\nu\alpha\beta}^{(a)}\bar{\Sigma}_{\lambda}^{\nu\alpha\dot{\alpha}}\Phi_{\dot{\alpha}}^{\mu\lambda(s)\beta}) \quad (84)$$

$$+12\sqrt{2}[\vec{O}_{(a)(R)}^{\mu\nu}\cdot(\vec{\Phi}_{\nu(R)}^{\lambda}\times\vec{\Sigma}_{\mu\lambda(R)}^{(s)})] \quad (85)$$

$$-\vec{O}_{\mu\nu(L)}^{(a)}\cdot(\vec{\Phi}_{\lambda(L)}^{\nu}\times\vec{\Sigma}_{(s)(L)}^{\mu\lambda})] \quad (86)$$

$$-6\sqrt{2}i(\Phi_{(a)}^{\nu\lambda\alpha\dot{\alpha}}\bar{\Sigma}_{\lambda\alpha}^{\mu\dot{\beta}}O_{\mu\nu\dot{\alpha}\dot{\beta}}^{(a)}+\Phi_{(a)}^{\nu\lambda\alpha\dot{\alpha}}\bar{\Sigma}_{\lambda\dot{\alpha}}^{\mu\beta}O_{\mu\nu\alpha\beta}^{(a)})] \quad (87)$$

The purely chiral superheavy supermultiplet masses can be determined from these expressions simply by substituting in the AM Higgs vevs and breaking up the contributions according to MSSM labels.

It is again easiest to keep track of Chiral fermion masses since all others follow using supersymmetry and the organization provided by the gauge super Higgs effect.

There are three types of mass terms involving fermions from chiral supermultiplets in such models :

- Unmixed Chiral
- Mixed pure chiral
- Mixed chiral-gauge. We briefly discuss the notable features of the mass spectrum calculation and give the actual mass formulae in the Appendix.

### 3.1 Unmixed Chiral

A pair of Chiral fermions transforming as  $SU(3) \times SU(2)_L \times U(1)_Y$  conjugates pairs up to form a massive Dirac fermion . For example for the properly normalized fields

$$\bar{A}[1, 1, -4] = \frac{\bar{\Sigma}_{44(R-)}}{\sqrt{2}} \quad A[1, 1, 4] = \frac{\Sigma_{44(R+)}}{\sqrt{2}} \quad (88)$$

one obtains the mass term

$$2(M + \eta(p + 3a + 6\omega))\bar{A}A = m_A\bar{A}A$$

The physical Dirac fermion mass is then  $|m_A|$  since the phase can be absorbed by a field redefinition. By supersymmetry this mass is shared by a pair of complex scalar fields with the same quantum numbers.

In the MSGUT case there are 19 pairs of chiral multiplets which form Dirac supermultiplets pairwise and two Majorana singletons, none of which mix with others of their ilk. In the NMSGUT 6 of these pairs become mixed with others of the same type leaving 11 Dirac supermultiplets and 2 Majorana supermultiplets (S,Q) which are unmixed. If the representation is real rather than complex one obtains an extra factor of 2 in the masses. The relevant representations, field components and masses are given in Table 1.

### 3.2 Mixed Pure Chiral

For such multiplets there is no mixing with the massive coset gauginos but there is a mixing among several multiplets with the same SM quantum numbers. There were only three such multiplet types in the MSGUT (i.e  $R[8, 1, 0]$ ,  $h[1, 2, \pm 1]$ ,  $t[3, 1, \pm \frac{2}{3}]$ ) but in the NMSGUT, as already mentioned there are an additional 5 mixed pure chiral

types namely the  $C[8, 2, \pm 1]$ ,  $D[3, 2, \pm \frac{7}{3}]$ ,  $K[3, 1, \pm \frac{8}{3}]$ ,  $L[6, 1, \pm \frac{2}{3}]$ ,  $P[3, 3, \pm \frac{2}{3}]$ . As for the multiplet types which had mixed pure chiral mass terms in the MSGUT, the type  $R[8, 1, 0]$  acquires no new partners and has an unchanged mass matrix since the **120** has no such submultiplets. However the other two mixed pure chiral multiplet types of the MSGUT do acquire new contributions :

- $[1, 2, -1](\bar{h}_1, \bar{h}_2, \bar{h}_3, \bar{h}_4, \bar{h}_5, \bar{h}_6) \oplus [1, 2, 1](h_1, h_2, h_3, h_4, h_5, h_6) \equiv (H_2^\alpha, \bar{\Sigma}_2^{(15)\alpha}, \Sigma_2^{(15)\alpha}, \frac{\Phi_{44}^{2\alpha}}{\sqrt{2}}, O_2^\alpha, O_2^{(15)\alpha}) \oplus (H_{\alpha i}, \bar{\Sigma}_{\alpha i}^{(15)}, \Sigma_{\alpha i}^{(15)}, \frac{\Phi_{\alpha}^{44i}}{\sqrt{2}}, O_{\alpha i}, O_{\alpha i}^{(15)})$

Here one gets additional an 2 rows and 2 columns relative to the MSGUT since the 120-plet contains two pairs of doublets with MSSM type Higgs doublet quantum numbers so that the mass matrix  $\mathcal{H}$  is  $6 \times 6$ . To keep one pair of light doublets in the low energy effective theory, it is necessary to fine tune one of the parameters of the superpotential (e.g  $M_H$ ) so that  $\text{Det}\mathcal{H} = 0$ . By extracting the null eigenvectors of  $\mathcal{H}^\dagger \mathcal{H}$  and  $\mathcal{H} \mathcal{H}^\dagger$  one can compute the composition of the light doublet pair in terms of the doublet fields in the full SO(10) GUT, and, in particular, we can find the proportions of the doublets coming from the **10**, **126**, **120** multiplets which couple to the matter sector (see Section 5 and Appendix C). This information is crucial for investigating whether a fit of the fermion data accomplished by using the generic form of the SO(10) fermion mass formulae is compatible with the dictates of the MSGUT. In fact precisely such considerations led [14, 15] to the conclusion that the MSGUT is Type I Seesaw dominated yet gives too small neutrino masses.

- $[\bar{3}, 1, \frac{2}{3}](\bar{t}_1, \bar{t}_2, \bar{t}_3, \bar{t}_4, \bar{t}_5, \bar{t}_6, \bar{t}_7) \oplus [3, 1, -\frac{2}{3}](t_1, t_2, t_3, t_4, t_5, t_6, t_7) \equiv (H^{\bar{\mu}4}, \bar{\Sigma}_{(a)}^{\bar{\mu}4}, \Sigma_{(a)}^{\bar{\mu}4}, \Sigma_{(R0)}^{\bar{\mu}4}, \Phi_{4(R+)}^{\bar{\mu}}, O_{\bar{\mu}4(s)}^{\bar{\mu}}, O_{(R0)}^{\bar{\mu}4}) \oplus (H_{\bar{\mu}4}, \bar{\Sigma}_{\bar{\mu}4(a)}, \Sigma_{\bar{\mu}4(a)}, \bar{\Sigma}_{\bar{\mu}4(R0)}, \Phi_{\bar{\mu}(R-)}^4, O_{\bar{\mu}4(s)}^4, O_{\bar{\mu}4(R0)}^4)$

With the contribution of the **120**-plet one gets two additional rows and columns and the dimension of  $[3, 1, \pm \frac{2}{3}]$  mass matrix  $\mathcal{T}$  becomes  $7 \times 7$ . These triplets and antitriplets participate in baryon violating process since the exchange of  $(t_1, t_2, t_4, t_6, t_7) \oplus (\bar{t}_1, \bar{t}_2, \bar{t}_6, \bar{t}_7)$  Higgsinos generates  $d = 5$  operators of type QQQL and  $\bar{l}\bar{u}\bar{u}\bar{d}$ . The strength of the operator is controlled by the inverse of the  $\bar{t} - t$  mass matrix  $\mathcal{T}$ .

### 3.3 Mixed Chiral-Gauge

Finally we come to the mixing matrices for the chiral modes that mix with the gauge particles as well as among themselves. There is no direct mixing between MSSM fields contained in 120-plet with gauge particles. However mixing *is* present via other MSSM submultiplets present in MSGUT Higgs fields which further mix with gauge fields. This occurs for all such multiplet types except  $G[1, 1, 0]$  and  $X[3, 2, \pm \frac{4}{3}]$  which are unchanged, while for  $E[3, 2, \pm \frac{1}{3}]$ ,  $F[1, 1, \pm 2]$ ,  $J[3, 1, \pm \frac{4}{3}]$  mass matrices acquire additional rows and columns. Thus



- $[\bar{3}, 2, -\frac{1}{3}](\bar{E}_1, \bar{E}_2, \bar{E}_3, \bar{E}_4, \bar{E}_5, \bar{E}_6) \oplus [3, 2, \frac{1}{3}](E_1, E_2, E_3, E_4, E_5, E_6)$   
 $\equiv (\Sigma_{4i}^{\bar{\mu}\alpha}, \bar{\Sigma}_{4i}^{\bar{\mu}\alpha}, \phi_{(s)2}^{\bar{\mu}4\alpha}, \phi_2^{(a)\bar{\mu}4\alpha}, \lambda_2^{\bar{\mu}4\alpha}, O_{4i}^{\bar{\sigma}\alpha}) \oplus (\bar{\Sigma}_{\bar{\mu}\alpha 2}^4, \Sigma_{\bar{\mu}\alpha 2}^4, \phi_{\bar{\mu}4\alpha i}^{(s)}, \phi_{\bar{\mu}4\alpha i}^{(a)}, \lambda_{\bar{\mu}\alpha i}, O_{\bar{\sigma}\alpha}^{4i})$

The  $6 \times 6$  mass matrix  $\mathcal{E}$  has the usual super-Higgs structure : complex conjugates of the 5th row and column (omitting the diagonal entry) furnish left and right null eigenvectors of the chiral  $5 \times 5$  submatrix  $\mathbf{E}$  obtained by omitting the fifth row and column.  $\mathcal{E}$  has non zero determinant although the determinant of  $\mathbf{E}$  vanishes.

- $[1, 1, -2](\bar{F}_1, \bar{F}_2, \bar{F}_3, \bar{F}_4) \oplus [1, 1, 2](F_1, F_2, F_3, F_4) \equiv (\bar{\Sigma}_{44(R0)}/\sqrt{2}, \Phi_{(R-)}^{(15)}, \lambda_{(R-)}, O_{44}/\sqrt{2}) \oplus (\Sigma_{(R0)}/\sqrt{2}, \Phi_{(R+)}^{(15)}, \lambda_{(R+)}, O^{44}/\sqrt{2})$

The  $4 \times 4$  mass matrix  $\mathcal{F}$  has the usual super-Higgs structure : complex conjugates of the 3th row and column (omitting the diagonal entry) furnish left and right null eigenvectors of the chiral  $3 \times 3$  submatrix  $\mathbf{F}$  obtained by omitting the third row and column.  $\mathcal{F}$  has non zero determinant although the determinant of  $\mathbf{F}$  vanishes.

- $[\bar{3}, 1, -\frac{4}{3}](\bar{J}_1, \bar{J}_2, \bar{J}_3, \bar{J}_4, \bar{J}_5) \oplus [3, 1, \frac{4}{3}](J_1, J_2, J_3, J_4, J_5)$   
 $\equiv (\Sigma_{(R-)}^{\bar{\mu}4}, \phi_4^{\bar{\mu}}, \phi_4^{\bar{\mu}(R0)}, \lambda_4^{\bar{\mu}}, O_{(R-)}^{\bar{\mu}4}) \oplus (\bar{\Sigma}_{\bar{\mu}4(R+)}^4, \phi_{\bar{\mu}}^4, \phi_{\bar{\mu}(R0)}^4, \lambda_{\bar{\mu}}^4, O_{\bar{\mu}4}^{(R+)})$

The  $5 \times 5$  mass matrix  $\mathcal{J}$  also has the super-Higgs structure : complex conjugates of the 4th row and column (omitting the diagonal entry) furnish left and right null eigenvectors of the chiral  $4 \times 4$  submatrix  $\mathbf{J}$  obtained by omitting the fourth row and column.  $\mathcal{J}$  has non zero determinant although the determinant of  $\mathbf{J}$  vanishes.

This concludes our description of the superheavy mass spectrum of the NMSGUT, explicit details are given in the Appendix.

## 4 RG Analysis

In [5, 14], for the case of the MSGUT, we exhibited plots of the threshold corrections  $(\Delta_G, \Delta_W, \Delta_X)$  to  $\alpha_G(M_X)^{-1}$ ,  $\sin^2\theta_W$  and  $\log_{10} M_X$  versus  $\xi$ . Note that we follow the approach of [32] in which  $M_X$  is taken to be the mass of the lightest gauge multiplet which mediates proton decay and not the point where the 3 MSSM gauge couplings cross. In [15] we synthesized the three batches of information corresponding to the three roots of the cubic equation(8) by exhibiting contour plots of  $\Delta_G, \Delta_W, \Delta_X$  on the  $x$ -plane at representative values of the other ('slow') parameters  $\lambda, \eta, \gamma, \bar{\gamma}$ . In this section we summarize information on the  $x$  values allowed by imposing plausible 'realistic' constraints on the magnitudes of the threshold corrections to the gauge couplings. We pay particular attention to the scenario[23] where  $M_X$  and with it all dangerous triplet masses are pushed above  $10^{16} GeV$ . To implement the consistency

requirements that the  $SO(10)$  theory remain perturbative after threshold and two loop corrections and, conversely, that  $\alpha_G$  not decrease so much as to invalidate the neglect of one-loop effects in the chiral couplings we impose an upper limit of 10 on the change in the magnitude of  $\alpha_G^{-1}$ . Similarly we estimate that uncertainty concerning the value of the supersymmetry breaking masses translates to an uncertainty of up to 10% in  $\sin^2 \theta_W(M_S)$ . Finally we expect that the mass of the lightest gauge mediating baryon decay bosons should not be lowered by more than one order of magnitude in order to respect the current bounds on  $d = 6$  mediated nucleon decay and not be raised by more than 2 orders of magnitude so that it remains well below the Planck scale. Thus we demand :

$$\begin{aligned} |\Delta_G| &\equiv |\Delta(\alpha_G^{-1}(M_X))| \leq 10 \\ 2 \geq \Delta_X &\equiv \Delta(\text{Log}_{10} M_X) \geq -1 \\ |\Delta_W| &\equiv |\Delta(\sin^2 \theta_W(M_S))| < .02 \end{aligned} \quad (89)$$

We find threshold corrections [5, 14]

$$\begin{aligned} \Delta^{(th)}(\text{Log}_{10} M_X) &= .0217 + .0167 \sum_{M'} (5\bar{b}'_1 + 3\bar{b}'_2 - 8\bar{b}'_3) \text{Log}_{10} \frac{M'}{M_X} \\ \Delta^{(th)}(\sin^2 \theta_W(M_S)) &= .00004 - .00024 \sum_{M'} (4\bar{b}'_1 - 9.6\bar{b}'_2 + 5.6\bar{b}'_3) \text{Log}_{10} \frac{M'}{M_X} \\ \Delta^{(th)}(\alpha_G^{-1}(M_X)) &= .1565 + .01832 \sum_{M'} (5\bar{b}'_1 + 3\bar{b}'_2 + 12\bar{b}'_3) \text{Log}_{10} \frac{M'}{M_X} \end{aligned} \quad (90)$$

Where  $\bar{b}'_i = 16\pi^2 b'_i$  are 1-loop  $\beta$  function coefficients ( $\beta_i = b_i g_i^3$ ) for multiplets with mass  $M'$ . These corrections, together with the two loop gauge corrections, modify the one loop values corresponding to the successful gauge unification of the MSSM, see [5, 14, 15] for details.

The parameter  $\xi = \lambda M / \eta m$  is the only numerical parameter that enters into the cubic eqn.(8) that determines the parameter  $x$  in terms of which all the superheavy vevs are given. *It is thus the most crucial determinant of the mass spectrum .*

The rest of the coupling parameters divide into “diagonal” ( $\lambda, \eta, \rho$ ) and “non-diagonal” ( $\gamma, \bar{\gamma}, \zeta, \bar{\zeta}, k$ ) couplings with the latter exerting a very minor influence on the unification parameters. We have therefore fixed the non diagonal parameters at representative values  $\sim 1$  throughout. The dependence of the threshold corrections on the “diagonal” couplings is also comparatively mild except when coherent e.g when many masses are lowered together leading to  $\alpha_G$  explosion,  $\text{Log} M_X$  collapse or large changes in  $\sin^2 \theta_W(M_S)$ . This happens when we lower these couplings too much. For lack of space and cogent reason to be exhaustive we give sample plots to summarize the most characteristic variation of the threshold effects over the complex  $x$ -plane.

A crucial point[14] is that the threshold corrections depend only on ratios of masses and are independent of the overall scale parameter which we choose to be  $m$ . Since  $M_X = 10^{16.25+\Delta_X} GeV$  it follows that

$$\begin{aligned}\Delta_X &= \Delta(\text{Log}_{10} \frac{M_X}{1 GeV}) \\ |m| &= 10^{16.25+\Delta_X} \frac{|\lambda|}{g\sqrt{4|\tilde{a} + \tilde{w}|^2 + 2|\tilde{p} + \tilde{\omega}|^2}} GeV\end{aligned}\tag{91}$$

It is thus clear that this factor will enter every superheavy mass so that they must all rise or fall in tandem with  $M_X$  i.e with  $\Delta_X$ .

In Figs. 3,4,5, for the case of the old MSGUT, we give examples of regions of the complex  $x$ -plane which are compatible with the unification constraints (89) for characteristic representative values of the diagonal couplings. As either of the couplings  $\lambda, \eta$  is lowered below magnitudes of order unity the region of the  $x$ -plane where the three unification constraints (89) are simultaneously satisfied changes in a characteristic way. The twin lobes (Fig. 3) in the first and fourth quadrants rotate towards the second and third quadrants respectively (Figs 3,4,5). The same behaviour is observed in the case of the NMSGUT but with the twist that the dependence on the parameter  $\rho$  can induce a considerable broadening of the allowed region when  $\rho$  is in a critical region near  $\rho_c = .15$  (Figs 6,7,8) and the other diagonal couplings are  $\sim 1$ .

An even more remarkable feature of both these sets of plots is that the allowed regions of the  $x$ -plane are dominantly those where the unification scale is raised above  $10^{17.25} GeV$  (the darkest shaded parts of Figs. 3-8). Thus the NMSGUT seems to point unequivocally and generically towards a resolution of the difficulties with  $d = 5$  baryon decay and a too low gauge Landau pole by an across the board elevation of GUT scale masses.

As discussed above, due to the successful fit of fermion masses and CP violation using only real superpotential couplings, the fitting problem for the real case is even more interesting than the general case and certainly more tractable. If the superpotential parameters are real so is  $\xi$  and it follows that the 3 solutions  $x_i$  of eqn.(8) form a conjugate pair accompanied by a real solution or else are all independently real. We shall show in section 5 that spontaneous CP violation requires that  $x$  be complex. Due to the presence of a reflection symmetry that interchanges the complex conjugate pair of solutions of eqn.(8) it is sufficient to study solutions with positive real imaginary part only. The complex conjugate pair of solutions exists only for  $\xi > -27.916$  and are present as a conjugate pair. Then we may ask for what (real) values of  $\xi$  can we obtain complex  $x$  values compatible with the constraints of unification. As already seen the answer depends on the values of  $\lambda, \eta, \rho$ . In Fig. 9 we show a parametric plot (vs  $\xi$ ) of the branch  $x_+(\xi)$  of the solution of eqn.(8) with positive




Figure 3: MSGUT all couplings  $\sim 1$  : Regions of the  $x$ -plane compatible with the unification constraints (89) are shaded. The darkest regions have  $2 \geq \Delta_X > 1$  (corresponding to  $M_X > 10^{17.25} \text{GeV}$ ), the next darkest  $1 \geq \Delta_X > 0$  the lightest shade  $0 \geq \Delta_X > -1$  and the white regions are disallowed.

imaginary part. The reflection symmetry in the  $Re[x]$  axis makes discussion of just the positive imaginary part branch sufficient.

In Figs. 10,11,12 we exhibit 3D plots showing regions of parameter space where viable complex  $x$  values can be generated from real values of  $\xi$ . This is a first step in the goal of determining which regions of the parameter space could actually support the successful generic fits of the fermion data discussed above. It is remarkable that the combined requirements of spontaneous CP violation and unification leave only rather restricted regions of the diagonal parameter space viable. In particular




Figure 4: MSGUT with  $\lambda$  lowered: Regions of the  $x$ -plane compatible with the unification constraints (89) are shaded. The darkest regions have  $2 \geq \Delta_X > 1$  (corresponding to  $M_X > 10^{17.25} \text{ GeV}$ ), the next darkest  $1 \geq \Delta_X > 0$  the lightest shade  $0 \geq \Delta_X > -1$  and the white regions are disallowed.

small values of the diagonal parameters  $\lambda$  or  $\eta$  are not allowed : as may be seen by superposing Fig.9 on Figs.4,5 (which are for the MSGUT but are essentially similar to the NMSGUT plots for small  $\lambda, \eta$ ). The reason is that lowering  $\lambda, \eta$  results in lower thresholds and explosive growth of the gauge coupling. When both  $\lambda, \eta$  are  $\sim 1$ , however, one finds that the solution curve shown in Fig. 9 intersects the allowed region for  $\xi$  values from about  $\xi \sim 10$  to as low as  $\xi \sim -27.9$  (one can approach even closer to  $\xi \simeq -27.917$  below which all solutions are real by tuning  $\rho$  to its critical value).




Figure 5: MSGUT with  $\eta$  lowered : Regions of the  $x$ -plane compatible with the unification constraints (89) are shaded. Further lowering  $\eta$  gives a plot almost identical to Fig. 4 where the lowering is due to  $\lambda$ .

#### 4.1 Suppression of Baryon decay

As discussed in the introduction, a very interesting question is whether there are viable regions of the parameter space where the theory is still perturbative (i.e.  $\alpha_G^{-1}(M_X) \geq 10$ , the change in  $\sin^2 \theta_W$  is acceptable and  $M_X \ll M_{Planck}$ ) yet the masses of the colour triplet Higgsinos that mediate proton decay are sufficiently large as to remove or mitigate the challenge to GUTs posed by the non-detection of Baryon decay. Surprisingly-in view of the tight upper bound on the triplet masses in the renormalizable  $SU(5)$  theory[21] the answer is *generically* in the affirmative. The rule of thumb of [21] indicates that  $M_{Triplet} \geq 7 \times 10^{16} GeV$  is sufficient to sup-




Figure 6: NMSGUT critical behaviour :  $\rho = .01$  i.e much smaller than  $\rho_c = .15$ , all other couplings  $\sim 1$ . Regions of the  $x$ -plane compatible with the unification constraints (89) are shaded and are essentially like the MSGUT case. The darkest regions have  $2 \geq \Delta_X > 1$  (corresponding to  $M_X > 10^{17.25} GeV$ ), the next darkest  $1 \geq \Delta_X > 0$  the lightest shade  $0 \geq \Delta_X > -1$  and the white regions are disallowed.

press  $d = 5$  Baryon decay operators to acceptable levels. In our case (see Section 6) there is not one pair but a plethora of triplets -of three different MSSM types ( $t[3, 1, \pm \frac{2}{3}]$ ,  $P[3, 3, \pm \frac{2}{3}]$ ,  $K[3, 1, \pm \frac{8}{3}]$ )- that can mediate Baryon decay. We can use the smallest mass of each of these three types as a surrogate for the the SU(5)  $M_{Triplet}$ . However in view of the several multiplets involved it is clear that there will be additional mixing angles associated with the diagonalization of triplet mass matrices that will enter the baryon decay amplitudes. Hence the lower bound on  $M_{triplet}$  is likely




Figure 7: NMSGUT critical behaviour:  $\rho = \rho_c = .15$ . Regions of the  $x$ -plane compatible with the unification constraints (89) are shaded, note the broadening of the allowed region with merger of the allowed ‘lobes’.

to be somewhat weaker than quoted (say  $10^{16} GeV$  for definiteness).

In Fig. 13,14,15 we plot the values of  $\Delta_X, \Delta_G, \Delta_W$  versus  $\xi$  with other couplings chosen at a representative points in the viable ranges indicated by Figs 10-12 and with  $x = x_+(\xi)$  (for spontaneous CP violation). There is a sharp peak in  $m$  (the overall mass scale in all superheavy masses written in terms of the dimensionless vevs) due to a peak in  $\Delta_X$  at  $\xi = -5$ . (which on the  $x_+(\xi)$  branch corresponds to  $x = 1 + i$ ). On this complex branch this spike is not due to any special gauge symmetry as can be checked from the values of the  $SO(10)/G_{123}$  coset gaugino (i.e E,X,G,F,J type






Figure 8: NMSGUT critical behaviour :  $\rho = .51$  i.e larger than  $\rho_c = .15$  : Regions of the x-plane compatible with the unification constraints (89) are shaded, note the reversion to the two lobe pattern of the MSGUT.

gauginos ) masses which remain distinct and non zero. Rather it is because a certain special multiplet, namely the lowest mass eigenstate in the  $C[8, 2, \pm 1]$  sector becomes light as one approaches close to  $\xi = -5$  ( i.e as  $M_X, \alpha_G$  rise). One cannot approach the exact value too closely as  $\alpha_G$  explodes. This is visible in Fig. 16 where we show the Mass of this light state versus  $\xi$ . Note however that the ‘crevasse’ into which  $M_C^{min}$  is to fall is very narrow indeed and that outside the region  $\xi \in (-5.5, -4.5)$  both  $\alpha_G$  and  $M_C$  are comfortably normal. In fact, except exactly at the singularity  $M_C$  is always above  $10^{15} GeV$  since it too has a factor of  $m$  within it. The quantum numbers of this multiplet are reminiscent of the intermediate scale colour octet and




Figure 9: Parametric plot of  $x(\xi)$   $|Im[x(\xi)] > 0, \xi \in [-27.917, 1000)$ . The terminus point near  $(2.3, 0)$  corresponds to  $\xi \rightarrow -27.917$  and that near  $(1, 0)$  to  $\xi \rightarrow \infty$

$SU(2)$  triplet multiplets used in raising the unification mass in [23]. Note however that we follow the approach the effective field theory based approach of [32] who defines the unification scale  $M_X$  simply as the mass of the lightest B violating Gauge boson. It would be interesting to reanalyze these questions using other approaches to the RG flow. The multiplet  $C[8, 2, 1\pm]$  seems to have an analogous effect in this approach to that of the triplet and octet thresholds used in [23]. That is to say although its mass is still quite large outside the region  $\xi \in (-5.5, -4.5)$  where  $\alpha_G$  and  $M_X$  explode, yet it is already lowered sufficiently to raise  $M_X$  sufficiently for our purposes, as may be seen from Fig.13.  $\Delta_X$  never drops below about +.2, for  $\xi < 0$  it is always  $> 1.2$  and goes as high as 5.2 when one approaches  $\xi = -5$ . Apart from a limited special region around  $\xi \simeq 4$ ,  $\Delta_X$  is seen to exceed 1 over the entire  $\xi$  line. One unsatisfactory aspect of this approach is that although consistent with MSSM unification at  $10^{16.25} GeV$  it fails to justify it as a physical gauge threshold since even the lightest gauge field is heavier than the point where the couplings in the effective theory cross in the one loop theory.

In Table I we give a list of masses and parameter values for which  $M_X$  is raised to  $10^{18} GeV$ , while the gauge coupling of the spontaneously broken  $SO(10)$  theory is still quite small and we have used a value of  $\xi = -8.5$  which is far from the singularity. In fact approaching closer pushes  $M_X$  above  $10^{18} GeV$ .




Figure 10: Values of  $\lambda, \eta, \xi' = .02\xi$  compatible with unification stability constraints at  $\rho = 0.95$ .

The composition of the special  $C[8, 2, \pm 1]$  multiplet is worth mentioning. At the singular point it is a mixture of only the modes  $C_{1,2}, \bar{C}_{1,2}$  which have their origin in the  $(15, 2, 2)_{PS}$  submultiplets of the **126**,  $\overline{\mathbf{126}}$  multiplets. Thus they couple[5] to the matter fields as

$$W = 8\sqrt{2}f_{AB}(\bar{C}_1\bar{d}_AQ_B - C_2\bar{u}_AQ_B) + \dots \quad (92)$$

It is not clear to us if this rather banal coupling is of any phenomenological importance.

Even if one excludes the region close to  $\xi = -5$  in order to maintain perturbativity and to keep  $M_X < 10^{18} GeV$  (indicated by the horizontal line at  $\Delta_X = 1.75$  in Fig.13)

| Field[ $SU(3), SU(2), Y$ ]                           | $Masses(Units\ of\ 10^{18}Gev)$                                                                                                                                   |
|------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $A[1, 1, 4]$                                         | 1.19                                                                                                                                                              |
| $B[6, 2, 5/3]$                                       | 0.49                                                                                                                                                              |
| $C[8, 2, 1]$                                         | 1.13, 0.91, 0.27                                                                                                                                                  |
| $D[3, 2, 7/3]$                                       | 2.03, 1.74, 0.47                                                                                                                                                  |
| $E[3, 2, 1/3]$                                       | 2.12, 1.56, 1.01, 1.01, 0.78, 0.55                                                                                                                                |
| $F[1, 1, 2]$                                         | 2.16, 0.41, 0.41, 0.14                                                                                                                                            |
| $G[1, 1, 0]$                                         | 1.19, 0.85, 0.39, 0.33, 0.33, 0.06                                                                                                                                |
| $h[1, 2, 1]$                                         | 3.35, 2.46, 2.06, 1.34, 0.36                                                                                                                                      |
| $I[3, 1, 10/3]$                                      | 1.46                                                                                                                                                              |
| $J[3, 1, 4/3]$                                       | 1.99, 1.22, 0.61, 0.61, 0.22                                                                                                                                      |
| $K[3, 1, 8/3]$                                       | 1.23, 0.49                                                                                                                                                        |
| $L[6, 1, 2/3]$                                       | 1.45, 0.12                                                                                                                                                        |
| $M[6, 1, 8/3]$                                       | 1.44                                                                                                                                                              |
| $N[6, 1, 4/3]$                                       | 1.52                                                                                                                                                              |
| $O[1, 3, 2]$                                         | 3.86                                                                                                                                                              |
| $P[3, 3, 2/3]$                                       | 3.0, 0.21                                                                                                                                                         |
| $Q[8, 3, 0]$                                         | 1.29                                                                                                                                                              |
| $R[8, 1, 0]$                                         | 2.39, 0.59                                                                                                                                                        |
| $S[1, 3, 0]$                                         | 2.64                                                                                                                                                              |
| $t[3, 1, 2/3]$                                       | 3.47, 2.26, 1.44, 1.03, 0.73, 0.18, 0.12                                                                                                                          |
| $U[3, 3, 4/3]$                                       | 2.18                                                                                                                                                              |
| $V[1, 2, 3]$                                         | 1.69                                                                                                                                                              |
| $W[6, 3, 2/3]$                                       | 2.05                                                                                                                                                              |
| $X[3, 2, 5/3]$                                       | 0.97, 0.97, 0.6                                                                                                                                                   |
| $Y[6, 2, 1/3]$                                       | 0.7                                                                                                                                                               |
| $Z[8, 1, 2]$                                         | 2.37                                                                                                                                                              |
| <i>ParameterValues</i>                               | $\lambda = 1.0, \eta = .95, \gamma = .7, \bar{\gamma} = .6, \zeta = .7$<br>$\bar{\zeta} = .85, k = .35, m_0 = .8, \rho = .95, m/\lambda = 5.28 \times 10^{16}Gev$ |
| <i>Min[Mass](Units of <math>10^{18}Gev</math>)</i>   | 0.06( $G[1, 1, 0]$ ), 0.12( $L[6, 1, 2/3], t[3, 1, 2/3]$ )                                                                                                        |
| <i>Max[Mass](Units of <math>10^{18}Gev</math>)</i>   | 3.86( $O[1, 3, 2]$ )                                                                                                                                              |
| <i>Threshold Corrections</i>                         | $\Delta_X = 1.739, \Delta_W = -0.149, \Delta_G = 0.4994$<br>$\alpha_G = 0.038$                                                                                    |
| <i>GaugeMasses(Units of <math>10^{18}Gev</math>)</i> | $m_{\lambda_E} = 1.01, m_{\lambda_F} = 0.41, m_{\lambda_G} = 0.67,$<br>$m_{\lambda_J} = 0.61, m_{\lambda_X} = 0.97$                                               |

Table 1: Example of Masses and couplings favouring high  $M_X$ ,  $\xi = -8.5$  on the  $x = x_+(\xi)$  branch for real couplings




Figure 11: Values of  $\lambda, \rho, \xi' = .02\xi$  compatible with unification stability constraints at  $\eta = 0.95$ .

it is clear that negative  $\xi$  alone raises the Unification scale by more than an order of magnitude. This implies that in a region near  $\xi = -5$  the complex solutions are compatible with perturbative unification yet raise the Unification scale to values as close as desired to  $M_{Planck}$ . To maintain perturbativity, and the Weinberg angle prediction and allow neglect of gravitational radiative corrections it is sufficient to demand  $\Delta_X < 2$  so that  $M_X < 2 \times 10^{18} GeV \sim 10^{-1} M_P$ . From Fig 14 we see that the value of  $\alpha_G$  rises as one raises  $M_X$  but this is true only in a very narrow region around  $\xi = 5$  which is in any case excluded by too large a value of  $M_X$ . Otherwise




Figure 12: Values of  $\rho, \eta, \xi' = .02\xi$  compatible with unification stability constraints at  $\lambda = 1.00$ .

the change in  $\alpha_G$  from the one loop value is quite minor so that perturbativity in the gauge coupling is not an issue until one goes very close to the Planck scale. Obviously the  $d = 6$  i.e gauge mediated baryon decay operators operators will be completely suppressed when  $M_X$  is raised towards the Planck scale (say  $10^{18} GeV$  ) which will raise the  $d = 6$  operator mediated lifetime by 8 orders of magnitude above the  $10^{36}$  years usually quoted for  $d = 6$  processes in supersymmetric GUTs. In Fig. 15 we observe that the change in  $\sin^2 \theta_W$  only restricts one from going too far from the




Figure 13: Plot of  $\Delta_X$  against  $\xi$  on the CP violating solution branch  $x_+(\xi)$  at representative allowed values of the diagonal parameters.

singularity!

To see what happens with  $d = 5$  baryon decay operators we show in Figs 17,18,19 the masses of the lightest Baryon decay triplets of each of the three independent types  $t, K, P$  (see Section 6).

The mass of the common B-decay triplet  $t[3, 1, \pm \frac{2}{3}]$  is seen to always lie above  $10^{16} GeV$  and is even higher for most of the  $\xi$  line. We emphasize that the effective mass is greater since there will be a rotation matrix element between the lightest eigenstate and the modes that couple to the baryon violating  $d = 5$  operators (see Section 6). The situation with the  $P[3, 3, \pm \frac{2}{3}]$  and  $K[3, 1, \pm \frac{8}{3}]$  type triplets is even better since they mostly lie above  $10^{17} GeV$  !!

There is however a price to pay for all this mass scale raising and it is enforced by the other arm of the Baryon violation- Lepton violation seesaw or balance that operates in SO(10) theories. Namely (Fig. 19) the  $\overline{126}$  vev also rises above  $6 \times 10^{16} GeV$  for most of the  $\xi$  line. If one wishes to have right handed neutrinos much lighter than the GUT scale this would indicate that solutions of the fermion mass fitting problem with very small values of the  $\overline{126}$  Yukawa couplings are preferable to those that rely on inordinately skewed values of the Higgs fractions. Interestingly the semi-realistic 3 generation fits we obtained by perturbing around the 2-3 generation




Figure 14: Plot of  $\Delta_G$  against  $\xi$  on the CP violating solution branch  $x_+(\xi)$  at representative allowed values of the diagonal parameters.

**10 – 120** indeed have very small  $f_{AB}$  [16, 17](see Section 5). On the other hand the values of  $f$  are somewhat larger ( $\sim .1$ ) in the case of the sample realistic fit obtained using a numerical fitting procedure[18]. This may be an argument in favour of realistic fits close to our semi-realistic ones (which can be searched for[35] by starting the downhill simplex procedure near the solutions we found[17]). If indeed the viable solution finally has high **126** vev accompanied by the very small **126** Yukawa coupling required by the Type I Seesaw and the **10, 120** dominant charged fermion fit, it will be yet another instance of the intricate balance of the Fermion mass hierarchy and its ouroborotic link to the structure of the apparently completely remote GUT scale symmetry breaking and threshold structure.

Thus we see that the NMSGUT deals with the difficulty regarding  $d = 5$  mediated baryon decay in a very natural and unforced way. No fine tuning of Yukawas or artificially arranged cancellation or any introduction of a plethora of uncontrollable non-renormalizable terms is required at all. The suppression is generic in the viable regions of the parameter space and is practically inescapable on the complex branches of the solution of eqn.(8) which are structurally imposed requirements when CP violation is spontaneous.






Figure 15: Plot of  $\Delta_W$  against  $\xi$  on the CP violating solution branch  $x_+(\xi)$  at representative allowed values of the diagonal parameters.

## 4.2 Capping the Landau Pole

Furthermore, raising  $M_X$  to values near the Planck scale alleviates an even grosser difficulty of the MSGUT one loop unification scenario with  $\mathbf{210} \oplus \mathbf{126} \oplus \overline{\mathbf{126}}$  Higgs system without threshold effects [22] : the presence of a Landau pole in the gauge coupling evolution at  $\Lambda_X \sim 5M_X$  i.e lying just above the perturbative unification scale but well below the Planck scale. The problem is only worsened by the introduction of the  $\mathbf{120}$ . If the Unification scale is raised close to the Planck scale by the threshold effects that we have calculated then the strongly coupled dynamics at  $\Lambda_X$  occurs at or close to the Planck scale itself. This strengthens the heuristic arguments [25] that envision a (calculable) UV condensation of coset gauginos in the supersymmetric GUT (along with chiral vevs related by the Konishi anomaly and therefore an associated induced gravity a la AdS-CFT [28] ) which drives the breaking of the GUT symmetry[25]. It is worth remarking that such a scenario naturally overcomes all the fundamental objections[27] that led to an abandonment of the induced gravity scenarios of the 1980s. Moreover since the cutoff of the perturbative theory is about an order of magnitude less than the scale of UV condensation and the Planck scale - which coincide, it is natural to surmise that the gauge strong coupling dynamics




Figure 16: Plot of  $M_C^{min}$  against  $\xi$  on the CP violating solution branch  $x_+(\xi)$  at representative allowed values of the diagonal parameters.

induces gravity characterized by  $M_P \sim \Lambda_X$ . In this picture the MSSM Grand Desert evolution finds  $SO(10)$  completion when it crosses the superheavy mass thresholds and then  $SO(10)$  quickly defines its own physical UV cutoff :  $\Lambda_X \sim 5M_X$ . A supersymmetric theory with a physical cutoff escapes the objections (raised on grounds of ambiguity of cutoff dependent contributions[27]) against gravity induced by gauge theory dynamics. The coincidence of the scale of condensation and the Planck scale is of course the nub of the matter. Previous attempts to construct induced gravity from asymptotically free theories had no plausible reason at all why a large Planck scale should be induced by gauge theory(e.g QCD or some variety of Technicolour) without any intrinsic truly large scale in it at all. Here however we are ‘gifted’ with coincident Planck and strong *supersymmetric* (therefore holomorphically controlled and calculable[25]) condensation scales with no extra assumptions at all. Combined with the enlistment[23] of the MSSM singlets internal to the GUT in the cause of specifying the softly broken supersymmetry low energy theory, the whole scenario sounds almost too much like a free lunch to be true !




Figure 17: Plot of  $M_t^{min}$ , the lightest triplet of MSSM type  $[3, 1, \pm \frac{2}{3}]$ , against  $\xi$  on the CP violating solution branch  $x_+(\xi)$  at representative allowed values of the diagonal parameters. Units of  $10^{16} GeV$ .

## 5 Fermion Mass Formulae

As in the case of the MSGUT one imposes the fine tuning condition  $Det\mathcal{H} = 0$  to keep a pair of Higgs doublets  $H_{(1)}, \bar{H}_{(1)}$  (left and right null eigenstates of the mass matrix  $\mathcal{H}$ ) light. The composition of these null eigenstates in terms of the GUT scale doublets then specifies how much the different doublets contribute to the low energy EW scale symmetry breaking. In the Dirac mass matrices we can replace  $\langle h_i \rangle \rightarrow \alpha_i v_u, \langle \bar{h}_i \rangle \rightarrow \bar{\alpha}_i v_d$ . The fermion Dirac masses may be read off the decomposition of  $\mathbf{16} \cdot \mathbf{16} \cdot (\mathbf{10} \oplus \mathbf{120} \oplus \overline{\mathbf{126}})$  given in [3, 5] and this yields[15]

$$\begin{aligned}
 m^u &= v(\hat{h} + \hat{f} + \hat{g}) \quad ; \quad r_1 = \frac{\bar{\alpha}_1}{\alpha_1} \cot \beta \quad ; \quad r_2 = \frac{\bar{\alpha}_2}{\alpha_2} \cot \beta \\
 m^\nu &= v(\hat{h} - 3\hat{f} + (r_5 - 3)\hat{g}) \quad ; \quad r_5 = \frac{4i\sqrt{3}\alpha_5}{\alpha_6 + i\sqrt{3}\alpha_5} \\
 m^d &= v(r_1\hat{h} + r_2\hat{f} + r_6\hat{g}); \quad r_6 = \frac{\bar{\alpha}_6 + i\sqrt{3}\bar{\alpha}_5}{\alpha_6 + i\sqrt{3}\alpha_5} \cot \beta
 \end{aligned}$$




Figure 18: Plot of  $M_P^{min}$ , the lightest triplet of MSSM type  $[3, 3, \pm \frac{2}{3}]$ , against  $\xi$  on the CP violating solution branch  $x_+(\xi)$  at representative allowed values of the diagonal parameters. Units of  $10^{16} GeV$ .

$$\begin{aligned}
 m^l &= v(r_1 \hat{h} - 3r_2 \hat{f} + (\bar{r}_5 - 3r_6) \hat{g}); \quad \bar{r}_5 = \frac{4i\sqrt{3}\bar{\alpha}_5}{\alpha_6 + i\sqrt{3}\alpha_5} \cot \beta \\
 \hat{g} &= 2ig\sqrt{\frac{2}{3}}(\alpha_6 + i\sqrt{3}\alpha_5) \sin \beta \quad ; \quad \hat{h} = 2\sqrt{2}h\alpha_1 \sin \beta \quad ; \quad \hat{f} = -4\sqrt{\frac{2}{3}}if\alpha_2 \sin \beta
 \end{aligned} \tag{93}$$

The Yukawa couplings of matter fields with **120** Higgs field give no contribution to the Majorana mass matrix of the superheavy neutrinos  $\bar{\nu}_A$  so it remains  $M_{AB}^{\bar{\nu}} = 8\sqrt{2}f_{AB}\bar{\sigma}$ . Thus the Type I contribution is obtained by eliminating  $\bar{\nu}_A$

$$\begin{aligned}
 W &= \frac{1}{2}M_{AB}^{\bar{\nu}}\bar{\nu}_A\bar{\nu}_B + \bar{\nu}_A m_{AB}^{\nu} \nu_B + \dots \rightarrow \frac{1}{2}M_{AB}^{\nu(I)} \nu_A \nu_B + \dots \\
 M_{AB}^{\nu(I)} &= -((m^{\nu})^T (M^{\bar{\nu}})^{-1} m^{\nu})_{AB}
 \end{aligned} \tag{94}$$

As shown in [14, 15] it is likely that the Type II seesaw contribution is subdominant to the Type I seesaw. However the consistency of the assumption that it is negligible must be checked and quantified so we also evaluate the tadpole that gives rise to the Type II seesaw since the **120** plet does contribute new terms.




Figure 19: Plot of  $M_K^{min}$ , the lightest triplet of MSSM type  $[3, 1, \pm \frac{8}{3}]$ , against  $\xi$  on the CP violating solution branch  $x_+(\xi)$  at representative allowed values of the diagonal parameters. Units of  $10^{16} GeV$ .

For computing  $\overline{O}(\overline{10}, 3, 1)_{\overline{126}}$  vev, inspection of the mass spectrum (Appendix A) yields the relevant terms in the superpotential as

$$\begin{aligned}
W_{FM}^{\overline{\Sigma}} &= M_O \vec{\bar{O}} \cdot \vec{O} - \frac{\bar{\gamma}}{\sqrt{2}} H^{\alpha\dot{\alpha}} \Phi_{44\dot{\alpha}}^{\beta} \bar{O}_{\alpha\beta} - \frac{\gamma}{\sqrt{2}} H^{\alpha\dot{\alpha}} \Phi_{\dot{\alpha}}^{44\beta} O_{\alpha\beta} \\
&- 2\sqrt{2}i\eta(\Sigma_4^{4\alpha\dot{\alpha}} \Phi_{44\dot{\alpha}}^{\beta} \bar{O}_{\alpha\beta} + \bar{\Sigma}_4^{4\alpha\dot{\alpha}} \Phi_{\dot{\alpha}}^{44\beta} O_{\alpha\beta}) \\
&+ \bar{\zeta}[\frac{1}{2}O_{\dot{\alpha}}^{\beta}\Sigma_{44\alpha\beta}^{\bar{\Phi}}\bar{\Phi}^{44\alpha\dot{\alpha}} + O_4^{4\alpha\dot{\alpha}}\Sigma_{44\alpha\beta}^{\bar{\Phi}}\bar{\Phi}_{\dot{\alpha}}^{44\beta} \\
&+ \frac{1}{2}O_{\dot{\alpha}}^{\beta}\bar{\Sigma}_{\alpha\beta}^{44}\Phi_{44}^{\alpha\dot{\alpha}} - O_4^{4\alpha\dot{\alpha}}\bar{\Sigma}_{\alpha\beta}^{44}\Phi_{44\dot{\alpha}}^{\beta}] \\
&= M_O \bar{O}_- O_+ \bar{O}_- (\frac{i\bar{\gamma}}{\sqrt{2}}\bar{\alpha}_1 + 2i\sqrt{3}\eta\bar{\alpha}_3 + \sqrt{3}\bar{\zeta}\bar{\alpha}_6 + i\bar{\zeta}\bar{\alpha}_5)\bar{\alpha}_4 v_d^2 \\
&- O_+ (\frac{i\gamma}{\sqrt{2}}\alpha_1 + 2i\sqrt{3}\eta\alpha_2 - \sqrt{3}\zeta\alpha_6 + i\zeta\alpha_5)\alpha_4 v_u^2
\end{aligned} \tag{95}$$

So




Figure 20: Plot of the vev  $\bar{\sigma}$  of the  $\overline{126}$  field responsible for the righthanded neutrino mass, against  $\xi$  on the CP violating solution branch  $x_+(\xi)$  at representative allowed values of the diagonal parameters. Units of  $10^{16}GeV$ .

$$\langle \bar{O}_- \rangle = \left( \frac{i\gamma}{\sqrt{2}}\alpha_1 + 2i\sqrt{3}\eta\alpha_2 - \sqrt{3}\zeta\alpha_6 + i\zeta\alpha_5 \right) \alpha_4 \left( \frac{v_u^2}{M_O} \right) \quad (96)$$

and  $M_O$  can be read off from Table I to be  $M_O = 2(M + \eta(3a - p))$ . The Type II neutrino mass is then simply  $M_{AB}^\nu = 16if_{AB} < \bar{O}_- >$ .

### 5.1 Spontaneous CP violation and the Higgs Fractions $\alpha_i, \bar{\alpha}_i$

As is apparent from the fermion mass formulae the coefficients  $\alpha_i, \bar{\alpha}_i$  are critical for the phenomenology of these models. They are calculated by determining the null left and right eigenvectors of  $\mathcal{H}$  (the  $[1, 2, \pm 1]$  mass matrix) and can be used to verify the compatibility of the NMSGUT with the realistic generic fits now achievable[16, 17, 18]. To this end we list explicit expressions for the  $\alpha_i, \bar{\alpha}_i$  in Appendix C. An immediate application is to check the conditions under which fits of the fermion data like the spontaneous CP violating generic fit of [18] can be realized in the NMSGUT. Intuitively it is evident that complex GUT scale vevs should require complex values of  $x$ . One finds that the six independent phases[18] that appear in the generic spontaneous

CP violation case in **10** – **120** –  $\overline{\mathbf{126}}$  are given in terms of our quantities by (in the convention where  $\alpha_1 = \bar{\alpha}_1$  are both real so that the contributions of the **10**plet to all Dirac masses are real and the phase of  $\bar{\sigma}$  in the Type I seesaw formula has been absorbed by redefining the neutrino fields(as also in [18]) ):

$$\begin{aligned}
\zeta_u &= \text{Arg}[\alpha_2] - \frac{\pi}{2} \quad ; \quad \xi_u = \text{Arg}[i\alpha_6 - \sqrt{3}\alpha_5] \\
\zeta_d - \zeta_u &= \text{Arg}\left[\frac{\bar{\alpha}_2}{\alpha_2}\right] \quad ; \quad \xi_d - \xi_u = \text{Arg}\left[\frac{\bar{\alpha}_6 + i\sqrt{3}\bar{\alpha}_5}{\alpha_6 + i\sqrt{3}\alpha_5}\right] \\
\xi_l - \xi_u &= \text{Arg}[r_7] = \text{Arg}\left[\frac{-3\bar{\alpha}_6 + i\sqrt{3}\bar{\alpha}_5}{\alpha_6 + i\sqrt{3}\alpha_5}\right] \quad ; \quad \xi_D - \xi_u = \text{Arg}\left[\frac{-3\alpha_6 + i\sqrt{3}\alpha_5}{\alpha_6 + i\sqrt{3}\alpha_5}\right]
\end{aligned} \tag{97}$$

From Appendix C one can check that for real superpotential parameters and real values of  $x$  the  $\alpha_i, \bar{\alpha}_i, i = 1...5$  are real while  $\alpha_6, \bar{\alpha}_6$  are pure imaginary. Then it immediately follows that except the trivial  $\overline{\mathbf{126}}$  phase convention dependent phase values  $\zeta_u = \zeta_d = -\frac{\pi}{2}$  (essentially from the factor of  $i$  that accompanies  $f_{AB}$  within the parameter  $\hat{f}_{AB}$ ) all other phases are zero and there is no CKM CP violation. Clearly the only way to get non trivial phases in the CKM matrix while retaining real superpotential parameters is for  $x$  to be complex.

The generic fits **10** – **120** –  $\overline{\mathbf{126}}$  fits to the fermion data are to be carried out using eqns.(93) and the Type I seesaw masses [16, 17, 18]. They thus yield information only on the ratios

$$\begin{aligned}
\frac{\bar{\alpha}_1}{\alpha_1} &= r_1 \tan \beta \quad ; \quad \frac{\bar{\alpha}_2}{\alpha_2} = r_2 \tan \beta \\
\frac{\alpha_5}{\alpha_6} &= \frac{ir_5}{\sqrt{3}(r_5 - 4)} \\
\frac{\bar{\alpha}_5}{\alpha_6} &= \frac{i\bar{r}_5}{\sqrt{3}(r_5 - 4)} \tan \beta \\
\frac{\bar{\alpha}_6}{\alpha_6} &= \frac{(\bar{r}_5 - 4r_6)}{(r_5 - 4)} \tan \beta
\end{aligned} \tag{98}$$

The fitting procedure reduces our ignorance concerning the embedding of the MSSM in the MSGUT considerably and thus the stage is set by the formulae and analysis given here and in[15, 16, 17, 18] for better informed estimates of the exotic processes such as Baryon violation and Lepton flavour violation. The latter may be crucial for falsifiability tests if the high- $M_X$  route to suppress B-decay(and solve several other core problems !) is adopted. The critical unknowns are now the Leptonic CP violating phases but the theory is already sufficiently well specified as to permit a survey of the possibilities [35].

## 5.2 Sample fits

To illustrate the kind of realistic fit obtainable using the **10** – **120** –  **$\overline{126}$**  Higgs system we translate the  $\tan\beta = 10$  fit obtained in [18] into our notation and also give our own semi-realistic fits [17] for comparison. These expressions also allow us to easily see the special features required to ensure the consistency of the fit in terms of their implications for the  $\alpha_i, \bar{\alpha}_i$ . We emphasize however that these fits are significant purely as proof of principle. It may also be that the fits with much smaller values of  **$\overline{126}$**  Yukawas as found by us [17] may be preferred phenomenologically since they allow lowering of the righthanded neutrino mass even when the  **$\overline{126}$**  vev has been raised along with  $M_X$ . Relatively low right handed neutrino masses are often required required by Leptogenesis scenarios. This possible dovetailing of the requirements of baryon decay(suppression) and (left and right) neutrino phenomenology is perhaps another tantalizing glimpse of the ouroborotic balance [16] visible in the construction of MSGUTs.

## 5.3 Grimus and Kühbeck realistic fit

$$\begin{aligned}
\hat{h} &= \text{Diag}\{.000375, -.021266, .58331\} \\
\hat{f} &= e^{-0.0517654i} \begin{pmatrix} -0.00016545 & 0 & 0.00727765 \\ 0 & 0.0210717 & 0 \\ 0.00727765 & 0 & -0.105497 \end{pmatrix} \\
\hat{g} &= e^{3.95591i} \begin{pmatrix} 0 & .00031077 & 0 \\ -.00031077 & 0 & 0.0015122 \\ 0 & -0.0015122 & 0 \end{pmatrix} \\
r_1 &= .0109799 \quad ; \quad r_2 = 0.00309973 + 0.00129256i \\
r_5 &= 291.223 - 306.971i \quad ; \quad \bar{r}_5 = 0.19026 - 0.134289i \\
r_6 &= 0.110976 - 0.0852565i
\end{aligned} \tag{99}$$

From these one easily deduces

$$\begin{aligned}
\frac{\bar{\alpha}_1}{\alpha_1} &= r_1 \tan\beta = 0.1098 \quad ; \quad \frac{\bar{\alpha}_2}{\alpha_2} = r_2 \tan\beta = 0.03358e^{.395i} \\
\frac{\alpha_5}{\alpha_6} &= \frac{ir_5}{\sqrt{3}(r_5 - 4)} = e^{1.578i}.58112 \\
\frac{\bar{\alpha}_5}{\alpha_6} &= \frac{i\bar{r}_5 \tan\beta}{\sqrt{3}(r_5 - 4)} = e^{1.7748i}.003198 \\
\frac{\bar{\alpha}_6}{\alpha_6} &= \frac{(\bar{r}_5 - 4r_6) \tan\beta}{(r_5 - 4)} = e^{-1.436i}.00778
\end{aligned} \tag{100}$$



In [18] the vev ( $w_R$ ) of the  $\overline{\mathbf{126}}$  is taken somewhat arbitrarily to coincide with the GUT scale  $2 \times 10^{16}$  GeV. This could lead to inconsistencies since there is absolutely no reason why  $w_R$  can be assumed to take the fixed value  $2 \times 10^{16} \text{GeV}$ .

In [18] the increase in the size of neutrino masses is traced to the fact that  $f^{-1}$  is large and that a certain coefficient  $r_D$  has a large magnitude :  $|r_D| = 3008.9 \sim |(r_5 - 3)/r_6|$ . Since the denominator contains the two small numbers  $\frac{\bar{\alpha}_5}{\alpha_6}, \frac{\bar{\alpha}_6}{\alpha_6}$  we see that the largeness of  $r_D$  arises from the requirement that the fraction of the MSSM doublet  $\overline{H}$  contributed by the  $\mathbf{120}$  plet is very small. The same is evidently also true for the  $\overline{\mathbf{126}}$  since  $\frac{\bar{\alpha}_2}{\alpha_2}$  is also very small. It remains to be seen if these requirements are all in fact realizable in the NMSGUT or whether a different type of fit such as the one with much smaller  $\overline{\mathbf{126}}$  Yukawas which leads to very different values of the ratios of  $\alpha$ 's is more successful.

The fit of [18] was accomplished by fixing  $f_{12} = f_{23} = g_{13} = 0$  by imposing a  $Z_2$  symmetry and also by assuming the vev of the  $\overline{\mathbf{126}}$  to be  $M_X = 2 \times 10^{16}$  GeV. Since this vev enters only in the overall size of the neutrino masses it is clear that some leeway regarding that overall scale is still present if this assumption is not made given that they used the remaining parameters to fit the measured mass squared splittings of the neutrinos rather than the masses themselves. Moreover in the fit of [18] the leptonic CP phases emerge as (forced) outputs and have the values :

$$\delta^P = .21 \quad ; \quad \alpha_1^P = -3.1239 \quad ; \quad \alpha_2^P = -3.1404 \quad (101)$$

Since these phases have not yet been measured, and there no reason to assume that the  $Z_2$  symmetry and the assumption regarding the  $\overline{\mathbf{126}}$  vev have any basis in fact, it follows that we should regard these 4 parameters i.e  $f_{12}, f_{23}, g_{13}, \bar{\sigma}/M_X$  as lying in reserve for the day when the Leptonic CP angles and the neutrino mass scale are finally measured. This equality of the number of remaining unknowns and the number of remaining free parameters marks a rather poignant conjuncture for the NMSGUT leaving it poised on a razor's edge between inconsistency with the data and a grand leap towards an actual prediction of the rate of exotic processes such as baryon decay based firmly on an extrapolation of measured data and clear theoretical assumptions.

## 5.4 Semi-Realistic fits with ultra small $\overline{\mathbf{126}}$ Yukawas

$$\begin{aligned} \hat{r}_1 &= .016372 \quad ; \quad \hat{r}_2 = 0.98039 \quad ; \quad r_6 = 0.0187 \\ r_5 &= 2.67 \quad ; \quad \bar{r}_5 = 0.079405 \end{aligned}$$

$$\hat{h} = \begin{pmatrix} -0.0002988 & 0.0000394 & -0.0222691 \\ 0.0000394 & 0.009291 & -0.134823 \\ -0.0222691 & -0.134823 & 0.481333 \end{pmatrix}$$

$$\hat{f} = \begin{pmatrix} 0.0000132 & 0.0000277 & -0.0001 \\ 0.0000277 & -0.0000146 & 0.000057 \\ -0.0001 & 0.000057 & 0.000040 \end{pmatrix} \quad (102)$$

$$\hat{g} = \begin{pmatrix} 0 & 0.001644 & -0.02478 \\ -0.00164 & 0. & -0.119710 \\ 0.02478 & 0.119710 & 0 \end{pmatrix} \quad (103)$$

The eigenvalues of  $\hat{h}, \hat{f}, \hat{g}$  are

$$\begin{aligned} \hat{h} &: .518 \quad ; \quad .0277 \quad ; \quad 1.92 \times 10^{-5} \\ \hat{f} &: 1.33 \times 10^{-4} \quad ; \quad 1.11 \times 10^{-4} \quad ; \quad 0.171 \times 10^{-4} \\ \hat{g} &: \pm 0.122 \quad ; \quad 0 \end{aligned} \quad (104)$$

Observe that the values in  $f_{AB}$  are much smaller than the fit of [18]. It is also easy to check that the associated  $\alpha_i, \bar{\alpha}_i$  values are very different. It remains to be seen which type -if either- is actually realizable in the fully specified NMSGUT that we have described in this paper.

## 6 $d = 5$ Operators for B, L violation

In [5] we worked out the effective  $d = 4$  superpotential for  $B + L$  violating processes due to exchange of colour triplet superheavy chiral supermultiplets contained in the **10**,  **$\overline{126}$**  Higgs multiplets. These included a novel channel due to decays mediated by exchange of triplets  $t_{(4)}$  contained in the  **$\overline{126}$**  Higgs irrep. Evidently the inclusion of **120**plet Higgs will lead to additional channels for baryon violation. These can be easily derived using the Pati-Salam decomposition of the **16.16.120** SO(10) invariants[3] :

$$\begin{aligned} \frac{1}{(3!)} \psi C_2^{(5)} \gamma_i \gamma_j \gamma_k \chi O_{ijk} &= -2(\bar{O}_{(s)}^{\mu\nu} \psi_\mu^\alpha \chi_{\nu\alpha} + O_{\mu\nu}^{(s)} \hat{\psi}^{\mu\dot{\alpha}} \hat{\chi}_{\dot{\alpha}}^\nu) - 2\sqrt{2} O_\nu^{\mu\alpha\dot{\alpha}} (\hat{\psi}_{\dot{\alpha}}^\nu \chi_{\mu\alpha} - \psi_{\mu\alpha} \hat{\chi}_{\dot{\alpha}}^\nu) \\ &- 2(O_{\mu\nu}^{(a)\dot{\alpha}\dot{\beta}} \hat{\psi}_{\dot{\alpha}}^\mu \hat{\chi}_{\dot{\beta}}^\nu + \tilde{O}_{(a)}^{\mu\nu\alpha\beta} \psi_{\mu\alpha} \chi_{\nu\beta}) + \sqrt{2} O^{\alpha\dot{\alpha}} (\psi_{\dot{\alpha}}^\mu \chi_{\mu\alpha} - \psi_{\mu\alpha} \hat{\chi}_{\dot{\alpha}}^\mu) \end{aligned}$$

$$\begin{aligned} W_{FM}^O &= 2\sqrt{2} g_{AB} [\bar{h}_5 (\bar{d}_A \hat{Q}_B + \bar{e}_A \hat{L}_B) - h_5 (\hat{u}_A \hat{Q}_B + \hat{\nu}_A \hat{L}_B)] \\ &- 2\sqrt{2} g_{AB} [\sqrt{2} \bar{L}_2 \cdot (\hat{Q}_A \hat{Q}_B)' + F_4 \hat{L}_A \hat{L}_B + \sqrt{2} \bar{t}_6 \hat{Q}_A \hat{L}_B \\ &+ 2\sqrt{2} \bar{L}_2 \cdot (\hat{u}_A \bar{d}_B)' + \sqrt{2} t_6 (\hat{u}_A \bar{e}_B - \bar{d}_A \hat{\nu}_B) + 2\bar{F}_4 \bar{\nu}_A \bar{e}_B] \\ &- 2\sqrt{2} g_{AB} [2\bar{C}_3 \bar{d}_A \hat{Q}_B - 2C_3 \hat{u}_A \hat{Q}_B + \frac{i}{\sqrt{3}} \bar{h}_6 (\bar{d}_A \hat{Q}_B - 3\bar{e}_A \hat{L}_B) \end{aligned} \quad (105)$$

$$\begin{aligned}
& -\frac{i}{\sqrt{3}}h_6(\bar{u}_A\hat{Q}_B - 3\bar{\nu}_A\hat{L}_B) + 2\bar{D}_3\bar{e}_A\hat{Q}_B - 2\bar{E}_6\bar{\nu}_A\hat{Q}_B \\
& + 2E_6\bar{d}_A\hat{L}_B - 2D_3\bar{u}_A\hat{L}_B] - 2i\sqrt{2}g_{AB}[\epsilon\bar{J}_5\bar{d}_A\bar{d}_B \\
& + 2K_2\bar{d}_A\bar{e}_B - \epsilon\bar{K}_2\bar{u}_A\bar{u}_B - 2J_5\bar{u}_A\bar{\nu}_B \\
& - \sqrt{2}\epsilon\bar{t}_7\bar{d}_A\bar{u}_B - \sqrt{2}t_7(\bar{d}_A\bar{\nu}_B - \bar{e}_A\bar{u}_B)] - 2g_{AB}[\epsilon P_2\hat{Q}_A\hat{Q}_B + 2\bar{P}_2\hat{Q}_A\hat{L}_B]
\end{aligned}$$

We have suppressed  $G_{321}$  indices and used a sub-multiplet naming convention specified in Section 2, the primes in the terms containing colour sextets ( $L, \bar{L}$ ) are explained in the caption to Table 1. The carets on the fields indicate that they are fields of the NMSGUT and are unitarily related to those of the MSSM( $M_X$ ).

In order that the exchange of a Higgsino that couples to matter with a given  $B + L$  lead to a  $B + L$  violating  $d = 5$  operator in the effective theory at sub GUT energies it is necessary that it have a nonzero contraction with a conjugate (MSSM) representation Higgsino that couples to a matter chiral bilinear with a  $B + L$  different from the conjugate of the first  $B + L$  value. On inspection one finds that not only the familiar triplet types  $[\bar{3}, 1, \pm\frac{2}{3}] \subset \mathbf{120}$  i.e  $\{\bar{t}_{(6)}, \bar{t}_{(7)}\}[\bar{3}, 1, \frac{2}{3}]$  and  $\{t_{(6)}, t_{(7)}\}$  but also the novel exchange modes from the  $P[3, 3, \pm\frac{2}{3}]$  and  $K[3, 1, \pm\frac{8}{3}]$  multiplet types can contribute to baryon violation. In the case of the  $\overline{\mathbf{126}}$  the  $\bar{P}_1, \bar{K}_1 \subset \overline{\mathbf{126}}$  multiplets did couple to the fermions but  $P_1, \bar{K}_1 \subset \mathbf{126}$  did not. The  $\mathbf{120}$  however contains both  $P_2, \bar{P}_2$  and  $K_2, \bar{K}_2$ . Since these mix with  $P_1, \bar{P}_1$  and  $K_1, \bar{K}_1$ , a number of fresh contributions appear.

The multiplets  $P_2[3, 3, -\frac{2}{3}], \bar{P}_2[\bar{3}, 3, \frac{2}{3}], K_2[3, 1, -\frac{8}{3}], \bar{K}_2[\bar{3}, 1, \frac{8}{3}]$  satisfy the requirement regarding  $B + L$  quantum numbers of the fields they couple to. Note in particular that these novel exchanges always lead to contributions in which at least one and possibly both pairs of final state family indices are antisymmetrized.

On integrating out the heavy triplet Higgs supermultiplets one obtains the following additional effective  $d = 4$  Superpotential for Baryon Number violating processes in the NMSGUT to leading order in  $m_W/M_X$ . We have taken the opportunity to insert a missing overall sign and correct minor sub/super-script typos in [5] :

$$W_{eff}^{\Delta B \neq 0} = -\hat{L}_{ABCD}(\frac{1}{2}\epsilon\hat{Q}_A\hat{Q}_B\hat{Q}_C\hat{L}_D) - \hat{R}_{ABCD}(\epsilon\bar{e}_A\bar{u}_B\bar{u}_C\bar{d}_D) \quad (106)$$

where the coefficients are

$$\begin{aligned}
\hat{L}_{ABCD} &= \mathcal{S}_1^1\tilde{h}_{AB}\tilde{h}_{CD} + \mathcal{S}_1^2\tilde{h}_{AB}\tilde{f}_{CD} + \mathcal{S}_2^1\tilde{f}_{AB}\tilde{h}_{CD} + \mathcal{S}_2^2\tilde{f}_{AB}\tilde{f}_{CD} \\
&- \mathcal{S}_1^6\tilde{h}_{AB}\tilde{g}_{CD} - \mathcal{S}_2^6\tilde{f}_{AB}\tilde{g}_{CD} + \sqrt{2}(\mathcal{P}^{-1})_2^1\tilde{g}_{AC}\tilde{f}_{BD} \\
&- (\mathcal{P}^{-1})_2^2\tilde{g}_{AC}\tilde{g}_{BD}
\end{aligned} \quad (107)$$

and

$$\hat{R}_{ABCD} = \mathcal{S}_1^1\tilde{h}_{AB}\tilde{h}_{CD} - \mathcal{S}_1^2\tilde{h}_{AB}\tilde{f}_{CD} - \mathcal{S}_2^1\tilde{f}_{AB}\tilde{h}_{CD} + \mathcal{S}_2^2\tilde{f}_{AB}\tilde{f}_{CD}$$

$$\begin{aligned}
& - i\sqrt{2}\mathcal{S}_4^1\tilde{f}_{AB}\tilde{h}_{CD} + i\sqrt{2}\mathcal{S}_4^2\tilde{f}_{AB}\tilde{f}_{CD} \\
& + \mathcal{S}_6^1\tilde{g}_{AB}\tilde{h}_{CD} - i\mathcal{S}_7^1\tilde{g}_{AB}\tilde{h}_{CD} - \mathcal{S}_6^2\tilde{g}_{AB}\tilde{f}_{CD} + i\mathcal{S}_7^2\tilde{g}_{AB}\tilde{f}_{CD} \\
& + i\mathcal{S}_1^7\tilde{h}_{AB}\tilde{g}_{CD} - i\mathcal{S}_2^7\tilde{f}_{AB}\tilde{g}_{CD} + \sqrt{2}\mathcal{S}_4^7\tilde{f}_{AB}\tilde{g}_{CD} \\
& + i\mathcal{S}_6^7\tilde{g}_{AB}\tilde{g}_{CD} + \mathcal{S}_7^7\tilde{g}_{AB}\tilde{g}_{CD} - \sqrt{2}(\mathcal{K}^{-1})_1^2\tilde{f}_{AD}\tilde{g}_{BC} \\
& - (\mathcal{K}^{-1})_2^2\tilde{g}_{AD}\tilde{g}_{BC}
\end{aligned} \tag{108}$$

here  $\mathcal{S} = \mathcal{T}^{-1}$  and  $\mathcal{T}$  is the mass matrix for  $[3, 1, \pm 2/3]$ -sector triplets :  $W = \bar{t}^i \mathcal{T}_i^j t_j + \dots$ , while

$$\tilde{h}_{AB} = 2\sqrt{2}h_{AB} \quad \tilde{f}_{AB} = 4\sqrt{2}f_{AB} \quad \tilde{g}_{AB} = 4g_{AB} \tag{109}$$

The fields in terms of which we actually require the decay operators are those of the MSSM which do not carry carets and are unitarily related to the GUT fields[14]:

$$\begin{aligned}
\hat{Q}_A &= \mathcal{Q}_{A'A}^* Q_{A'} \quad ; \quad \hat{L}_A = \mathcal{L}_{A'A}^* L_{A'} \quad ; \quad \hat{u}_A = (V^u)_{A'A}^* \bar{u}_{A'} \\
\hat{d}_A &= (V^d)_{A'A}^* \bar{d}_{A'} \quad ; \quad \hat{e}_A = V_{A'A}^{e*} \bar{e}_{A'}
\end{aligned} \tag{110}$$

The Unitary  $MSSM \subset MSGUT$  embedding matrices  $\mathcal{Q}, L, V^{u,d,e}$  are fixed partly by the unitary ambiguity in the GUT fields which allows one to change  $h, f, g \rightarrow U^T(h, f, g)U$  at will, and partly by fitting the fermion mass data to the corresponding GUT expressions[14, 15, 16, 17]. This fitting procedure determines the embedding matrices up to certain phases and furnishes expressions for the GUT couplings in terms of the fermion mass data of the MSSM( $M_X$ ) and the embedding matrices. So far solutions of the highly non-linear fitting problem have been found either by expanding around the dominant 2-3 sector of the fermion hierarchy[15, 16, 17] or by making ad-hoc simplifying assumptions concerning the Yukawa textures to reduce the number of unknowns and then using a non-linear fitting method such as the “downhill simplex” [18]. The complicating factor is that we still lack some important information on the overall mass scale of neutrinos, leptonic mixing ( $\theta_{13}^P$ ) and leptonic CP phase information  $\delta^P, \alpha_1^P, \alpha_2^P$ . Till they are available one can at most survey the possibilities[35]

## 7 Discussion and Outlook

In this paper, motivated by successful fits of the fermion data[17, 18] which evade the difficulties that forced an abandonment[14, 15] of the hope[9] that the **10** ,  **$\overline{126}$**  FM Higgs system would be sufficient to describe the entire fermion mass spectrum, we specified the ingredients of a New Minimal Supersymmetric GUT based on the gauge group  $SO(10)$  and the  **$210 \oplus 10 \oplus 120 \oplus 126 \oplus \overline{126}$**  Higgs System. While inheriting the Higgs system responsible for GUT scale symmetry breaking unchanged from

the MSGUT[10, 11, 12], and fitting all the fermion data successfully, the NMSGUT is able to describe CKM CP violation[18] even when it carries the handicap of real superpotential couplings. Since this handicap reduces the number of free parameters drastically it results in a theory with one less free superpotential parameter than even the original MSGUT! Moreover several parameters are yet available to accommodate the new information associated with the as yet unknown Leptonic CP violation phases,  $\theta_{13}^P$  angle and the over all neutrino mass scale. If this information became available it would determine the unknown but crucial embedding matrices describing the relation between the effective MSSM in the grand desert and the NMSGUT.

Using the techniques we developed for the MSGUT[3, 5] we computed the super-heavy spectrum for the NMSGUT and used it to compute threshold effects in the gauge evolution. Focussing on the symmetry breaking branch which allows spontaneous CP violation we found that the Unification scale defined as the mass of the Baryon number violating gauge fields was always raised above the one loop values on this branch. This increase could take  $M_X$  to values as large as  $10^{18} \text{ GeV}$  while still remaining in the perturbative domain. Thus gauge mediated Baryon decay can easily become unmeasurably small in this theory. Together with  $M_X$  all other masses, in particular those of the three triplet types that mediate  $d = 5$  baryon decay also rise and can be taken(effectively) well above  $10^{16} \text{ GeV}$ . Thus not only  $d = 6$  but also  $d = 5$  baryon decay- which has persistently been a hobgoblin in the fairy story of Grand Unification- can be banished to the periphery -if not completely outside -of the measurable realm.

Increase in  $M_X$  to  $\sim 10^{18} \text{ GeV}$  required by casting the GUT field in the role of supersymmetry breaker ( *sugrone* ?) calls for a reanalysis of the spontaneous breaking and spectral determination using the full Sugra potential. Since the GUT singlets are well poised to take over the role normally played by the Polonyi singlet or the hidden sector [23], we expect the minimum of the global Susy theory to morph into a more rigorous solution which also incorporates supersymmetry breaking and suppresses the cosmological constant. The participation of singlets from the  $\overline{126}$  which couples directly to the matter multiplets implies that there may be some (small) deviation from universality in the soft supersymmetry breaking masses that arise.

The increase of  $M_X$  provides resolution of another nagging and persistent difficulty[22] in the MSGUT : the Landau pole in the gauge coupling evolution above  $M_X$ . Since  $M_X$  is closer to the Planck scale the presence of the SO(10) Landau pole next to the Planck scale only strengthens our speculation that the UV condensation to be expected in such a supersymmetric Asymptotically Strong(AS) theory [22, 25] acts as both a physical cutoff for the perturbative SO(10) theory and the scale of an induced gravity that arises from this theory. Much remains to be done to implement this scenario but we made a beginning in[25] by demonstrating using Supersymmetric strong coupling heuristics[26] that in a toy ASSGUT the condensation actually takes place and breaks the (toy) GUT symmetry and that the vevs responsible are *calculable*.

It is encouraging that the development of the theory in regard to apparently unrelated features has naturally brought us to the point where a number of intractable fundamental features have suddenly become pliable to a synthetic interpretation.

We also gave complete formulae for the fermion masses and baryon violating effective superpotential in the NMSGUT, including very lengthy analytic expressions for the Higgs fractions ( $\alpha_i, \bar{\alpha}_i$ ) which are determined by the GUT parameters (after a fine tuning) and are crucial ingredients of both the masses and the  $d = 5$  B-violation. We illustrated some aspects of their use when examining the realizability of a generic fit within the NMSGUT with real parameters by taking the proof of principle fit of [18] as an example. However we note again that much remains to be done to properly map out the space of fermion fits in which the fit of [18] represents only an isolated point of no particular sanctity. Indeed, if the semi-realistic fits we found in [17] using a quite different method based on an expansion around the 2-3 generation core of the fermion hierarchy extend to the realistic case with CP violation, we expect the new fits to be distinct from those found in [18] and have much smaller values of **126** couplings. This distinction would then have important phenomenological implications since it would lead to much lower right handed neutrino masses ; as indeed is required in many Leptogenesis scenarios.

In consequence of the lowered rates for  $d = 5$  processes which long represented the main hope for detection of Baryon decay and for falsifying Susy GUT models, it may not be possible to detect such decays, at least in the foreseeable future. In that case Lepton Flavour violation [34] will remain the only arena where this theory has predictions that may be testable in the near term.

In sum, the NMSGUT having inherited the strengths of its parent is revealing new virtues that illuminate what was mysterious in the MSGUT and promise to carry the slow caravan of Grand unification not only across the Grand Desert that was its first horizon but across threshold jungles beyond that first horizon up into the rarefied heights where gauge forces and gravity meld into their primordial pleromal [25] unity.

## 8 Acknowledgments

The work of C.S.A was supported by a grant No SR/S2/HEP-11/2005 from the Department of Science and Technology of the Govt. of India and that of S.K.G by a University Grants Commission Junior Research Fellowship.

| Field[ $SU(3), SU(2), Y$ ]                                    | $PS$                                                                                                               | $Fields$ | Mass                               |
|---------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------|----------|------------------------------------|
| $A[1, 1, 4], \bar{A}[1, 1, -4]$                               | $\frac{\Sigma_{(R+)}^{44}}{\sqrt{2}}, \frac{\bar{\Sigma}_{44(R-)}}{\sqrt{2}}$                                      |          | $2(M + \eta(p + 3a + 6\omega))$    |
| $M[6, 1, \frac{8}{3}], \bar{M}[(\bar{6}, 1, -\frac{8}{3})]$   | $(\bar{\Sigma}_{\bar{\mu}\bar{\nu}(R+)}^{(R+)}, \Sigma'_{(R-)}{}^{\bar{\mu}\bar{\nu}})_{\bar{\mu} \leq \bar{\nu}}$ |          | $2(M + \eta(p - a + 2\omega))$     |
| $N[6, 1, -\frac{4}{3}], \bar{N}[(\bar{6}, 1, \frac{4}{3})]$   | $(\bar{\Sigma}_{\bar{\mu}\bar{\nu}}^{(R-)}, \Sigma'_{(R+)}{}^{\bar{\mu}\bar{\nu}})_{\bar{\mu} \leq \bar{\nu}}$     |          | $2(M + \eta(p - a - 2\omega))$     |
| $O[1, 3, -2], \bar{O}[(1, 3, +2)]$                            | $\frac{\bar{\Sigma}_{44(L)}}{\sqrt{2}}, \frac{\bar{\Sigma}_{(L)}^{44}}{\sqrt{2}}$                                  |          | $2(M + \eta(3a - p))$              |
| $W[6, 3, \frac{2}{3}], \bar{W}[(\bar{6}, 3, -\frac{2}{3})]$   | $\bar{\Sigma}'_{\bar{\mu}\bar{\nu}(L)}, \bar{\Sigma}_{(L)}^{\bar{\mu}\bar{\nu}}$                                   |          | $2(M - \eta(a + p))$               |
| $I[3, 1, \frac{10}{3}], \bar{I}[(\bar{3}, 1, -\frac{10}{3})]$ | $\phi_{\bar{\nu}(R+)}^4, \phi_{4(R-)}^{\bar{\nu}}$                                                                 |          | $-2(m + \lambda(p + a + 4\omega))$ |
| $S[1, 3, 0]$                                                  | $\bar{\phi}_{(L)}^{(15)}$                                                                                          |          | $2(m + \lambda(2a - p))$           |
| $Q[8, 3, 0]$                                                  | $\bar{\phi}_{\bar{\mu}(L)}^{\bar{\nu}}$                                                                            |          | $2(m - \lambda(a + p))$            |
| $U[3, 3, \frac{4}{3}], \bar{U}[(\bar{3}, 3, -\frac{4}{3})]$   | $\bar{\phi}_{\bar{\mu}(L)}^4, \bar{\phi}_{4(L)}^{\bar{\mu}}$                                                       |          | $-2(m - \lambda(p - a))$           |
| $V[1, 2, -3], \bar{V}[1, 2, 3]$                               | $\frac{\phi_{4\alpha\dot{2}}}{\sqrt{2}}, \frac{\phi_{\alpha\dot{1}}^{44}}{\sqrt{2}}$                               |          | $2(m + 3\lambda(a + \omega))$      |
| $B[6, 2, \frac{5}{3}], \bar{B}[(\bar{6}, 2, -\frac{5}{3})]$   | $(\phi'_{\bar{\mu}\bar{\nu}\alpha\dot{1}}, \phi_{\alpha\dot{2}}^{\bar{\mu}\bar{\nu}})_{\bar{\mu} \leq \bar{\nu}}$  |          | $-2(m + \lambda(\omega - a))$      |
| $Y[6, 2, -\frac{1}{3}], \bar{Y}[(\bar{6}, 2, \frac{1}{3})]$   | $(\phi'_{\bar{\mu}\bar{\nu}\alpha\dot{2}}, \phi_{\alpha\dot{1}}^{\bar{\mu}\bar{\nu}})_{\bar{\mu} \leq \bar{\nu}}$  |          | $2(m - \lambda(a + \omega))$       |
| $Z[8, 1, 2], \bar{Z}[8, 1, -2]$                               | $\phi_{\bar{\mu}(R+)}^{\bar{\nu}}, \phi_{\bar{\mu}(R-)}^{\bar{\nu}}$                                               |          | $2(m + \lambda(p - a))$            |

Table 2: (i) Masses of the unmixed states in terms of the superheavy vevs . The  $SU(2)_L$  contraction order is always  $\bar{F}^\alpha F_\alpha$ . The absolute value of the expressions in the column “Mass” is understood. For sextets of  $SU(3)$  the 6 unit norm fields are denoted by a prime :  $\Sigma'_{\bar{\mu}\bar{\nu}} = \Sigma_{\bar{\mu}\bar{\nu}}, \bar{\mu} > \bar{\nu}, \Sigma'_{\bar{\mu}\bar{\mu}} = \Sigma_{\bar{\mu}\bar{\mu}}/\sqrt{2}$  and similarly for  $\bar{6}$ .

## Appendix A : Tables of masses and mixings

Here mixing matrix rows are labelled by barred irreps and columns by unbarred.

(i) The masses of 13 Unmixed cases are given as Table II.

ii) **Mixed states**

$$\mathbf{a)} \quad [8, 2, -1](\bar{C}_1, \bar{C}_2, \bar{C}_3) \oplus [8, 2, 1](C_1, C_2, C_3) \equiv (\bar{\Sigma}_2^{A\alpha}, \Sigma_2^{A\alpha}, O_2^{A\alpha}) \oplus (\Sigma_{\alpha\dot{1}}^A, \bar{\Sigma}_{\alpha\dot{1}}^A, O_{\alpha\dot{1}}^A) (A = 1, \dots, 8)$$

$$\begin{pmatrix} 2(-M + \eta(a + \omega)) & 0 & -i(\omega - p)\bar{\zeta} \\ 0 & 2(-M + \eta(a + \omega)) & -i(\omega + p)\zeta \\ i(\omega - p)\zeta & i(\omega + p)\bar{\zeta} & -m_0 + \frac{2}{3}a \end{pmatrix}$$

$$\text{b)} \quad [\bar{3}, 2, -\frac{7}{3}](\bar{D}_1, \bar{D}_2, \bar{D}_3) \oplus [3, 2, \frac{7}{3}](D_1, D_2, D_3) \equiv (\Sigma_{42}^{\bar{\nu}\alpha}, \bar{\Sigma}_{42}^{\bar{\nu}\alpha}, O_{42}^{\bar{\nu}\alpha}) \oplus (\bar{\Sigma}_{\bar{\nu}\alpha\bar{1}}^4, \Sigma_{\bar{\nu}\alpha\bar{1}}^4, O_{\bar{\nu}\alpha\bar{1}}^4)$$

$$\begin{pmatrix} 2(M + \eta(a + \omega)) & 0 & (i\omega + ip - 2ia)\zeta \\ 0 & 2(M + \eta(a + 3\omega)) & (-3i\omega - ip - 2ia)\bar{\zeta} \\ -(i\omega + ip - 2ia)\bar{\zeta} & (3i\omega + ip + 2ia)\zeta & m_0 + \frac{\rho}{3}(a + 2\omega) \end{pmatrix}$$

$$\text{c)} \quad [\bar{3}, 2, -\frac{1}{3}](\bar{E}_1, \bar{E}_2, \bar{E}_3, \bar{E}_4, \bar{E}_5, \bar{E}_6) \oplus [3, 2, \frac{1}{3}](E_1, E_2, E_3, E_4, E_5, E_6) \\ \equiv (\Sigma_{41}^{\bar{\mu}\alpha}, \bar{\Sigma}_{41}^{\bar{\mu}\alpha}, \phi_{(s)2}^{\bar{\mu}4\alpha}, \phi_2^{(a)\bar{\mu}4\alpha}, \lambda_2^{\bar{\mu}4\alpha}, O_{41}^{\bar{\sigma}\alpha}) \oplus (\bar{\Sigma}_{\bar{\mu}\alpha\bar{2}}^4, \Sigma_{\bar{\mu}\alpha\bar{2}}^4, \phi_{\bar{\mu}4\alpha\bar{1}}^{(s)}, \phi_{\bar{\mu}4\alpha\bar{1}}^{(a)}, \lambda_{\bar{\mu}\alpha\bar{1}}, O_{\bar{\sigma}\alpha}^{4\bar{1}})$$

$$\begin{pmatrix} -2(M + \eta(a - \omega)) & 0 & 0 & 0 & 0 & (i\omega - ip + 2ia)\zeta \\ 0 & -2(M + \eta(a - 3\omega)) & -2\sqrt{2}\eta\sigma & 2i\eta\sigma & ig\sqrt{2}\bar{\sigma}^* & (-3i\omega + ip + 2ia)\bar{\zeta} \\ 0 & 2i\sqrt{2}\eta\bar{\sigma} & -2(m + \lambda(a - \omega)) & -2\sqrt{2}\lambda\omega & 2g(a^* - \omega^*) & -\sqrt{2}\bar{\zeta}\bar{\sigma} \\ 0 & -2i\eta\bar{\sigma} & -2\sqrt{2}\lambda\omega & -2(m - \lambda\omega) & \sqrt{2}g(\omega^* - p^*) & \bar{\sigma}\bar{\zeta} \\ 0 & -ig\sqrt{2}\sigma^* & 2g(a^* - \omega^*) & g\sqrt{2}(\omega^* - p^*) & 0 & 0 \\ (-i\omega + ip - 2ia)\bar{\zeta} & (3i\omega - ip - 2ia)\zeta & -\sqrt{2}\zeta\sigma & \sigma\zeta & 0 & -(m_0 + \frac{\rho}{3}a - \frac{2}{3}\rho\omega) \end{pmatrix}$$

$$\text{d)} \quad [1, 1, -2](\bar{F}_1, \bar{F}_2, \bar{F}_3, \bar{F}_4) \oplus [1, 1, 2](F_1, F_2, F_3, F_4) \equiv (\frac{\bar{\Sigma}_{44(R0)}}{\sqrt{2}}, \Phi_{(R-)}^{(15)}, \lambda_{(R-)}, \frac{O_{44}}{\sqrt{2}}) \oplus (\frac{\Sigma_{(R0)}^{44}}{\sqrt{2}}, \Phi_{(R+)}^{(15)}, \lambda_{(R+)}, \frac{O_{44}}{\sqrt{2}})$$

$$\begin{pmatrix} 2(M + \eta(p + 3a)) & -2i\sqrt{3}\eta\sigma & -g\sqrt{2}\bar{\sigma}^* & -6i\bar{\zeta}\omega \\ 2i\sqrt{3}\eta\bar{\sigma} & 2(m + \lambda(p + 2a)) & \sqrt{24}ig\omega^* & \sqrt{3}\bar{\zeta}\bar{\sigma} \\ -g\sqrt{2}\sigma^* & -\sqrt{24}ig\omega^* & 0 & 0 \\ 6i\zeta\omega & \sqrt{3}\zeta\sigma & 0 & m_0 + a\rho \end{pmatrix}$$

$$\text{e)} \quad [1, 1, 0](G_1, G_2, G_3, G_4, G_5, G_6) \equiv (\phi, \phi^{(15)}, \phi_{(R0)}^{(15)}, \frac{\Sigma_{(R-)}^{44}}{\sqrt{2}}, \frac{\bar{\Sigma}_{44((R+))}}{\sqrt{2}}, \frac{\sqrt{2}\lambda^{(R0)} - \sqrt{3}\lambda^{(15)}}{\sqrt{5}})$$

$$\mathcal{G} = 2 \begin{pmatrix} m & 0 & \sqrt{6}\lambda\omega & \frac{i\eta\bar{\sigma}}{\sqrt{2}} & \frac{-i\eta\sigma}{\sqrt{2}} & 0 \\ 0 & m + 2\lambda a & 2\sqrt{2}\lambda\omega & i\eta\bar{\sigma}\sqrt{\frac{3}{2}} & -i\eta\sigma\sqrt{\frac{3}{2}} & 0 \\ \sqrt{6}\lambda\omega & 2\sqrt{2}\lambda\omega & m + \lambda(p + 2a) & -i\eta\sqrt{3}\bar{\sigma} & i\sqrt{3}\eta\sigma & 0 \\ \frac{i\eta\bar{\sigma}}{\sqrt{2}} & i\eta\bar{\sigma}\sqrt{\frac{3}{2}} & -i\eta\sqrt{3}\bar{\sigma} & 0 & M + \eta(p + 3a - 6\omega) & \frac{\sqrt{5}g\sigma^*}{2} \\ \frac{-i\eta\sigma}{\sqrt{2}} & -i\eta\sigma\sqrt{\frac{3}{2}} & i\eta\sqrt{3}\sigma & M + \eta(p + 3a - 6\omega) & 0 & \frac{\sqrt{5}g\bar{\sigma}^*}{2} \\ 0 & 0 & 0 & \frac{\sqrt{5}g\sigma^*}{2} & \frac{\sqrt{5}g\bar{\sigma}^*}{2} & 0 \end{pmatrix}$$

$$\text{f)} \quad [1, 2, -1](\bar{h}_1, \bar{h}_2, \bar{h}_3, \bar{h}_4, \bar{h}_5, \bar{h}_6) \oplus [1, 2, 1](h_1, h_2, h_3, h_4, h_5, h_6) \equiv (H_2^\alpha, \bar{\Sigma}_2^{(15)\alpha}, \Sigma_2^{(15)\alpha}, \frac{\Phi_{44}^{2\alpha}}{\sqrt{2}}, O_2^\alpha, O_2^{(15)\alpha}) \oplus (H_{\alpha\bar{1}}, \bar{\Sigma}_{\alpha\bar{1}}^{(15)}, \Sigma_{\alpha\bar{1}}^{(15)}, \frac{\Phi_{\alpha}^{44\bar{1}}}{\sqrt{2}}, O_{\alpha\bar{1}}, O_{\alpha\bar{1}}^{(15)})$$

$$\begin{pmatrix} -M_H & \bar{\gamma}\sqrt{3}(\omega - a) & -\gamma\sqrt{3}(\omega + a) & -\bar{\gamma}\bar{\sigma} & kp & -\sqrt{3}ik\omega \\ -\bar{\gamma}\sqrt{3}(\omega + a) & 0 & -(2M + 4\eta(a + \omega)) & 0 & -\sqrt{3}\bar{\zeta}\omega & i(p + 2\omega)\bar{\zeta} \\ \gamma\sqrt{3}(\omega - a) & -(2M + 4\eta(a - \omega)) & 0 & -2\eta\bar{\sigma}\sqrt{3} & \sqrt{3}\zeta\omega & -i(p - 2\omega)\zeta \\ -\sigma\gamma & -2\eta\sigma\sqrt{3} & 0 & -2m + 6\lambda(\omega - a) & \zeta\sigma & \sqrt{3}i\zeta\sigma \\ pk & \sqrt{3}\bar{\zeta}\omega & -\sqrt{3}\omega\zeta & \bar{\zeta}\bar{\sigma} & -m_o & \frac{\rho}{\sqrt{3}}i\omega \\ \sqrt{3}ik\omega & i(p - 2\omega)\bar{\zeta} & -i(p + 2\omega)\zeta & -\sqrt{3}i\bar{\zeta}\bar{\sigma} & -\frac{\rho}{\sqrt{3}}i\omega & -m_0 - \frac{2\rho}{3}a \end{pmatrix}$$



The above matrix is to be diagonalized after imposing the fine tuning condition  $Det\mathcal{H} = 0$  to keep one pair of doublets light.

$$\begin{aligned} \text{g)} \quad & [\bar{3}, 1, -\frac{4}{3}](\bar{J}_1, \bar{J}_2, \bar{J}_3, \bar{J}_4, \bar{J}_5) \oplus [3, 1, \frac{4}{3}](J_1, J_2, J_3, J_4, J_5) \\ & \equiv (\Sigma_{(R-)}^{\bar{\mu}4}, \phi_4^{\bar{\mu}}, \phi_4^{\bar{\mu}(R0)}, \lambda_4^{\bar{\mu}}, O_{(R-)}^{\bar{\mu}4}) \oplus (\bar{\Sigma}_{\bar{\mu}4(R+)}^{\bar{\mu}4}, \phi_{\bar{\mu}}^4, \phi_{\bar{\mu}(R0)}^4, \lambda_{\bar{\mu}}^4, O_{\bar{\mu}4}^{(R+)}) \end{aligned}$$

$$\mathcal{J} = \begin{pmatrix} 2(M + \eta(a + p - 2\omega)) & -2\eta\bar{\sigma} & 2\sqrt{2}\eta\bar{\sigma} & -ig\sqrt{2}\sigma^* & 2\zeta(a - 2\omega) \\ 2\eta\sigma & -2(m + \lambda a) & -2\sqrt{2}\lambda\omega & -2ig\sqrt{2}a^* & -\sigma\zeta \\ -2\sqrt{2}\eta\sigma & -2\sqrt{2}\lambda\omega & -2(m + \lambda(a + p)) & -4ig\omega^* & \sqrt{2}\sigma\zeta \\ -ig\sqrt{2}\bar{\sigma}^* & 2\sqrt{2}iga^* & 4ig\omega^* & 0 & 0 \\ 2\bar{\zeta}(a - 2\omega) & \bar{\sigma}\bar{\zeta} & -\sqrt{2}\bar{\sigma}\bar{\zeta} & 0 & m_o + \frac{\rho}{3}(p - 2\omega) \end{pmatrix}$$

$$\text{h)} \quad [\bar{3}, 1, \frac{8}{3}](\bar{K}_1, \bar{K}_2) \oplus [3, 1, -\frac{8}{3}](K_1, K_2) \equiv (\Sigma_{(R+)}^{\bar{\mu}4}, O_{(R+)}^{\bar{\mu}4}) \oplus (\bar{\Sigma}_{\bar{\mu}4(R-)}^{\bar{\mu}4}, O_{\bar{\mu}4(R-)}^{\bar{\mu}4})$$

$$\begin{pmatrix} 2(M + \eta(a + p + 2\omega)) & 2\zeta(a + 2\omega) \\ 2\bar{\zeta}(a + 2\omega) & m_0 + \frac{\rho}{3}(p + 2\omega) \end{pmatrix}$$

$$\begin{aligned} \text{i)} \quad & [\bar{6}, 1, -\frac{2}{3}](\bar{L}_1, \bar{L}_2) \oplus [6, 1, \frac{2}{3}](L_1, L_2) \equiv (\Sigma_{(R0)}'^{\bar{\mu}\bar{\nu}(s)}, O'^{\bar{\mu}\bar{\nu}(s)}) \oplus \\ & (\bar{\Sigma}_{\bar{\mu}\bar{\nu}(s)(R0)}', O_{\bar{\mu}\bar{\nu}(s)}') \end{aligned}$$

$$\begin{pmatrix} 2(M + \eta(p - a)) & -2i\zeta\omega \\ 2i\bar{\zeta}\omega & m_0 - \frac{\rho}{3}a \end{pmatrix}$$

$$\text{j)} \quad [\bar{3}, 3, \frac{2}{3}](\bar{P}_1, \bar{P}_2) \oplus [3, 3, -\frac{2}{3}](P_1, P_2) \equiv (\vec{\Sigma}_{(L)}^{\bar{\mu}4}, \vec{O}_{(L)}^{\bar{\mu}4}) \oplus (\vec{\Sigma}_{\bar{\mu}4(L)}^{\bar{\mu}4}, \vec{O}_{\bar{\mu}4(L)}^{\bar{\mu}4})$$

$$\begin{pmatrix} 2(M + \eta(a - p)) & 2a\bar{\zeta} \\ 2a\zeta & m_0 - \frac{\rho}{3}p \end{pmatrix}$$

$$\text{k)} \quad [8, 1, 0](R_1, R_2) \equiv (\hat{\phi}_{\bar{\mu}}^{\bar{\nu}}, \hat{\phi}_{\bar{\mu}(R0)}^{\bar{\nu}})$$

$$\mathcal{R} = 2 \begin{pmatrix} (m - \lambda a) & -\sqrt{2}\lambda\omega \\ -\sqrt{2}\lambda\omega & m + \lambda(p - a) \end{pmatrix}$$

$$\mathbf{l)} \quad [\bar{3}, 1, \frac{2}{3}](\bar{t}_1, \bar{t}_2, \bar{t}_3, \bar{t}_4, \bar{t}_5, \bar{t}_6, \bar{t}_7) \oplus [3, 1, -\frac{2}{3}](t_1, t_2, t_3, t_4, t_5, t_6, t_7) \equiv (H^{\bar{\mu}4}, \bar{\Sigma}_{(a)}^{\bar{\mu}4}, \Sigma_{(a)}^{\bar{\mu}4}, \Sigma_{(R0)}^{\bar{\mu}4}, \Phi_{4(R+)}^{\bar{\mu}}, O_{(R0)}^{\bar{\mu}4(s)}, O_{(R0)}^{\bar{\mu}4}) \oplus (H_{\bar{\mu}4}, \bar{\Sigma}_{\bar{\mu}4(a)}, \Sigma_{\bar{\mu}4(a)}, \bar{\Sigma}_{\bar{\mu}4(R0)}, \Phi_{\bar{\mu}(R-)}^4, O_{\bar{\mu}4(s)}, O_{\bar{\mu}4(R0)})$$

$$\begin{pmatrix} M_H & \bar{\gamma}(a+p) & \gamma(p-a) & 2\sqrt{2}i\omega\bar{\gamma} & i\bar{\sigma}\bar{\gamma} & \sqrt{2}ka & \sqrt{2}ik\omega \\ \bar{\gamma}(p-a) & 0 & 2M & 0 & 0 & \sqrt{2}a\bar{\zeta} & \sqrt{2}i\omega\bar{\zeta} \\ \gamma(p+a) & 2M & 0 & 4\sqrt{2}i\omega\eta & 2i\eta\bar{\sigma} & -\sqrt{2}a\zeta & \sqrt{2}i\omega\zeta \\ -2\sqrt{2}i\omega\gamma & -4\sqrt{2}i\omega\eta & 0 & 2M+2\eta p+2\eta a & -2\sqrt{2}\eta\bar{\sigma} & 2i\omega\zeta & 2\zeta a \\ i\sigma\gamma & 2i\eta\sigma & 0 & 2\sqrt{2}\eta\sigma & -2m-2\lambda(a+p-4\omega) & \sqrt{2}i\sigma\zeta & -\sqrt{2}\zeta\sigma \\ \sqrt{2}ka & -\sqrt{2}a\bar{\zeta} & \sqrt{2}a\zeta & -2i\bar{\zeta}\omega & \sqrt{2}i\bar{\zeta}\bar{\sigma} & m_0+\frac{\rho}{3}a & -\frac{2i}{3}\rho\omega \\ -\sqrt{2}ik\omega & -\sqrt{2}i\omega\bar{\zeta} & -\sqrt{2}i\omega\zeta & 2\zeta a & \sqrt{2}\bar{\sigma}\bar{\zeta} & \frac{2i}{3}\rho\omega & m_0+\frac{\rho}{3}p \end{pmatrix}$$

$$\mathbf{m)} \quad [3, 2, \frac{5}{3}](\bar{X}_1, \bar{X}_2, \bar{X}_3) \oplus [3, 2, -\frac{5}{3}](X_1, X_2, X_3) \\ \equiv (\phi_{\alpha\dot{1}}^{(s)\bar{\mu}4}, \phi_{\alpha\dot{1}}^{(a)\bar{\mu}4}, \lambda_{\alpha\dot{1}}^{\bar{\mu}4}) \oplus (\phi_{\bar{\mu}4\alpha\dot{2}}^{(s)}, \phi_{\bar{\mu}4\alpha\dot{2}}^{(a)}, \lambda_{\bar{\mu}4\alpha\dot{2}})$$

$$\mathcal{X} = \begin{pmatrix} 2(m+\lambda(a+\omega)) & -2\sqrt{2}\lambda\omega & -2g(a^*+\omega^*) \\ -2\sqrt{2}\lambda\omega & 2(m+\lambda\omega) & \sqrt{2}g(\omega^*+p^*) \\ -2g(a^*+\omega^*) & \sqrt{2}g(\omega^*+p^*) & 0 \end{pmatrix}$$

## Appendix B : SU(5) × U(1) Reassembly Crosscheck

The internal consistency of these spectra and couplings can be verified by considering special values of vevs, e.g

$$p = a = \pm\omega \quad (111)$$

where the unbroken symmetry includes SU(5)[4]. Then we find that the MSSM labelled mass spectra and couplings given in Appendix A do indeed reassemble into SU(5) invariant form. If we insert  $a = -\omega = p$  in the mass matrices of Appendix A we find that, after diagonalizing the mass matrices of the submultiplets that mix, the resultant spectra group precisely as indicated by the decompositions below with all the subreps of a given SU(5) irrep obtaining the same mass and correct phases to permit reassembly. The delicacy of this reassembly is a very non-trivial consistency check of our results.

$$\begin{aligned} \mathbf{H} &= 10 = 5_1 + \bar{5}_{-1} \\ 5_1 &= h_1(1, 2, 1) + t_1(3, 1, -\frac{2}{3}) \\ \bar{5}_{-1} &= \bar{h}_1(1, 2, -1) + \bar{t}_1(\bar{3}, 1, \frac{2}{3}) \end{aligned} \quad (112)$$

$$\begin{aligned}
\Sigma &= 126 = 1_{-5}(G_4) + \bar{5}_{-1} + 10_{-3} + \overline{15}_3 + 45_1 + \overline{50}_{-1} \\
\bar{5}_{-1} &= \bar{h}_3(1, 2, -1) + \bar{t}_{3,4}(\bar{3}, 1, \frac{2}{3}) \\
10_{-3} &= F_1(1, 1, 2) + \bar{J}_1(\bar{3}, 1, -\frac{4}{3}) + E_2(3, 2, \frac{1}{3}) \\
\overline{15}_3 &= O(1, 3, -2) + \bar{E}_1(\bar{3}, 2, -\frac{1}{3}) + \bar{N}(\bar{6}, 1, \frac{4}{3}) \\
45_1 &= h_3(1, 2, 1) + t_3(3, 1, -\frac{2}{3}) + P_1(3, 3, -\frac{2}{3}) + \bar{K}_1(\bar{3}, 1, \frac{8}{3}) + \bar{D}_1(\bar{3}, 2, -\frac{7}{3}) \\
&\quad + \bar{L}_1(\bar{6}, 1, -\frac{2}{3}) + C_1(8, 2, 1) \\
\overline{50}_{-1} &= A(1, 1, 4) + \bar{t}_{3,4}(\bar{3}, 1, \frac{2}{3}) + D_2(3, 2, \frac{7}{3}) + W(6, 3, \frac{2}{3}) + \overline{M}(\bar{6}, 1, -\frac{8}{3}) + \overline{C}_2(8, 2, -1)
\end{aligned} \tag{113}$$

$$\begin{aligned}
\overline{\Sigma} &= \overline{126} = 1_5(G_5) + 5_1 + \overline{10}_3 + 15_{-3} + \overline{45}_{-1} + 50_1 \\
5_1 &= h_2(1, 2, 1) + t_{2,4}(3, 1, -\frac{2}{3}) \\
\overline{10}_3 &= \bar{F}_1(1, 1, -2) + J_1(3, 1, \frac{4}{3}) + \bar{E}_2(\bar{3}, 2, -\frac{1}{3}) \\
15_{-3} &= \bar{O}(1, 3, 2) + E_1(3, 2, \frac{1}{3}) + N(6, 1, -\frac{4}{3}) \\
\overline{45}_{-1} &= \bar{h}_2(1, 2, -1) + \bar{t}_2(3, 1, \frac{2}{3}) + \bar{P}_1(\bar{3}, 3, \frac{2}{3}) + K_1(3, 1, -\frac{8}{3}) + D_1(3, 2, \frac{7}{3}) \\
&\quad + L_1(6, 1, \frac{2}{3}) + \bar{C}_1(8, 2, -1) \\
50_1 &= \bar{A}(1, 1, -4) + t_{2,4}(3, 1, -\frac{2}{3}) + \bar{D}_2(\bar{3}, 2, -\frac{7}{3}) + \overline{W}(\bar{6}, 3, -\frac{2}{3}) + M(6, 1, \frac{8}{3}) + C_2(8, 2, 1)
\end{aligned} \tag{114}$$

$$\begin{aligned}
\Phi &= 210 = 1_0 + 5_{-4} + \bar{5}_4 + 10_2 + \overline{10}_{-2} + 24_0 + 40_2 + \overline{40}_{-2} + 75_0 \\
1_0 &= G_{1,2,3} \\
5_{-4} &= h_4(1, 2, 1) + t_5(3, 1, -\frac{2}{3}) \\
\bar{5}_4 &= \bar{h}_4(1, 2, -1) + \bar{t}_5(\bar{3}, 1, \frac{2}{3}) \\
10_2 &= F_2(1, 1, 2) + \bar{J}_{2,3}(\bar{3}, 1, -\frac{4}{3}) + E_{3,4}(3, 2, \frac{1}{3}) \\
\overline{10}_{-2} &= \bar{F}_2(1, 1, -2) + J_{2,3}(3, 1, \frac{4}{3}) + \bar{E}_{3,4}(\bar{3}, 2, -\frac{1}{3}) \\
24_0 &= (1, 1, 0)G_{1,2,3} + S(1, 3, 0) + X_{1,2}(3, 2, -\frac{5}{3}) + \bar{X}_{1,2}(\bar{3}, 2, \frac{5}{3}) + R_{1,2}(8, 1, 0) \\
40_2 &= V(1, 2, -3) + E_{3,4}(3, 2, \frac{1}{3}) + \bar{J}_{2,3}(\bar{3}, 1, -\frac{4}{3}) + \bar{U}(\bar{3}, 3, -\frac{4}{3}) + Z(8, 1, 2) + \bar{Y}(\bar{6}, 2, \frac{1}{3})
\end{aligned}$$

$$\begin{aligned}
\overline{40}_{-2} &= \bar{V}(1, 2, 3) + \bar{E}_{3,4}(3, 2, -\frac{1}{3}) + J_{2,3}(3, 1, \frac{4}{3}) + U(3, 3, \frac{4}{3}) + \bar{Z}(8, 1, -2) + Y(6, 2, -\frac{1}{3}) \\
75 &= (1, 1, 0)G_{1,2,3} + I(3, 1, \frac{10}{3}) + \bar{I}(\bar{3}, 1, -\frac{10}{3}) + X_{1,2}(3, 2, -\frac{5}{3}) + \bar{X}_{1,2}(\bar{3}, 2, \frac{5}{3}) \\
&+ B(6, 2, \frac{5}{3}) + \bar{B}(\bar{6}, 2, -\frac{5}{3}) + R_{1,2}(8, 1, 0) + Q(8, 3, 0)
\end{aligned} \tag{115}$$

$$\begin{aligned}
\mathbf{O} &= 5_1 + \bar{5}_{-1} + 10_{-3} + \bar{10}_3 + 45_1 + \bar{45}_{-1} \\
5_1 &= h_{5,6}(1, 2, 1) + t_{6,7}(3, 1, -\frac{2}{3}) \\
\bar{5}_{-1} &= \bar{h}_{5,6}(1, 2, -1) + \bar{t}_{6,7}(\bar{3}, 1, \frac{2}{3}) \\
10_{-3} &= F_4(1, 1, 2) + \bar{J}_5(\bar{3}, 1, -\frac{4}{3}) + E_6(3, 2, \frac{1}{3}) \\
\bar{10}_3 &= \bar{F}_4(1, 1, -2) + J_5(3, 1, \frac{4}{3}) + \bar{E}_6(\bar{3}, 2, -\frac{1}{3}) \\
45_1 &= h_{5,6}(1, 2, 1) + t_{6,7}(3, 1, -\frac{2}{3}) + P_2(3, 3, -\frac{2}{3}) + \bar{K}_2(\bar{3}, 1, \frac{8}{3}) + \bar{D}_3(\bar{3}, 2, -\frac{7}{3}) \\
&+ \bar{L}_2(6, 1, -\frac{2}{3}) + C_3(8, 2, 1) \\
\bar{45}_{-1} &= \bar{h}_{5,6}(1, 2, -1) + \bar{t}_{6,7}(\bar{3}, 1, \frac{2}{3}) + \bar{P}_2(\bar{3}, 3, \frac{2}{3}) + K_2(3, 1, -\frac{8}{3}) + D_3(3, 2, \frac{7}{3}) \\
&+ L_2(6, 1, \frac{2}{3}) + \bar{C}_3(8, 2, -1)
\end{aligned} \tag{116}$$

Due to the **120**plet one obtains the additional  $SU(5)$  invariant mass terms:

$$\begin{aligned}
&(m_0 + \rho p)5_O \bar{5}_O + (m_0 + \rho p)10_O \bar{10}_O + (m_0 - \frac{\rho}{3}p)45_O \bar{45}_O \\
&+ 2kp (5_O \bar{5}_H + \bar{5}_O 5_H) - 2\sqrt{3}p(\zeta 5_O \bar{5}_\Sigma + \bar{\zeta} \bar{5}_O 5_\Sigma) + 2(\zeta \sigma 5_O \bar{5}_\Phi + \bar{\zeta} \bar{\sigma} \bar{5}_O 5_\Phi) \\
&+ 6ip(\bar{\zeta} 10_O \bar{10}_\Sigma + \zeta \bar{10}_O 10_\Sigma) + \sqrt{3}(\bar{\zeta} \bar{\sigma} 10_O \bar{10}_\Phi + \zeta \sigma \bar{10}_O 10_\Phi) \\
&+ 2p(\bar{\zeta} 45_O \bar{45}_\Sigma + \zeta \bar{45}_O 45_\Sigma)
\end{aligned} \tag{117}$$

Where every  $SU(5)$  invariant has been normalized so that the individual  $G_{123}$  sub-rep masses can be read off directly from the coefficient of the invariant for complex  $SU(5)$  representations which pair into Dirac supermultiplets and is 2 times the coefficient for the real representations which remain unpaired Majorana/Chiral supermultiplets.

### Appendix C : Doublet fraction Coefficients $\alpha_i, \bar{\alpha}_i$

In this appendix we give the explicit expressions for the coefficients  $\alpha_i, \bar{\alpha}_i$  obtained by first imposing the condition  $Det\mathcal{H} = 0$  and then solving the equations to determine the normalized left and right eigenvectors of  $\mathcal{H}$ .

$$\tilde{m}_0 = \frac{m_0 \lambda}{m} \quad \hat{\sigma} = \sqrt{\frac{(1-3x)x(1+x^2)}{(1-2x+x^2)}}$$

$$N = \frac{1}{\sqrt{|\hat{\alpha}_1|^2 + |\hat{\alpha}_2|^2 + |\hat{\alpha}_3|^2 + |\hat{\alpha}_4|^2 + |\hat{\alpha}_5|^2 + |\hat{\alpha}_6|^2}}$$

$$\bar{N} = \frac{1}{\sqrt{|\hat{\bar{\alpha}}_1|^2 + |\hat{\bar{\alpha}}_2|^2 + |\hat{\bar{\alpha}}_3|^2 + |\hat{\bar{\alpha}}_4|^2 + |\hat{\bar{\alpha}}_5|^2 + |\hat{\bar{\alpha}}_6|^2}}$$

$$A = N\{\hat{\alpha}_1, \hat{\alpha}_2, \hat{\alpha}_3, \hat{\alpha}_4, \hat{\alpha}_5, \hat{\alpha}_6\} = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6\}$$

$$\bar{A} = \bar{N}\{\hat{\bar{\alpha}}_1, \hat{\bar{\alpha}}_2, \hat{\bar{\alpha}}_3, \hat{\bar{\alpha}}_4, \hat{\bar{\alpha}}_5, \hat{\bar{\alpha}}_6\} = \{\bar{\alpha}_1, \bar{\alpha}_2, \bar{\alpha}_3, \bar{\alpha}_4, \bar{\alpha}_5, \bar{\alpha}_6\}$$

$$\hat{\alpha}_1 = \hat{\bar{\alpha}}_1 = (\tilde{m}_o^2 \eta^2 \lambda P_0 + \bar{\zeta}^2 \zeta^2 \lambda P_1 + \tilde{m}_o \bar{\zeta} \zeta \eta \lambda P_2 + \bar{\zeta} \zeta \eta \lambda \rho P_3 + \tilde{m}_o \eta^2 \lambda \rho P_4 + \eta^2 \lambda \rho^2 P_5)$$

$$\begin{aligned} \hat{\alpha}_2 = & (\tilde{m}_o^2 \gamma \eta \lambda Q_0 + \tilde{m}_o \bar{\zeta} \gamma \zeta \lambda Q_1 + \bar{\gamma} \tilde{m}_o \zeta^2 \lambda Q_2 + k \bar{\zeta} \zeta^2 \lambda Q_3 + \bar{\zeta}^2 \gamma \zeta^2 \lambda Q_4 + \bar{\gamma} \bar{\zeta} \zeta^3 \lambda Q_5 \\ & + k \tilde{m}_o \zeta \eta \lambda Q_6 + \bar{\zeta} \gamma \zeta \lambda \rho Q_7 + \bar{\gamma} \zeta^2 \lambda \rho Q_8 + \tilde{m}_o \gamma \eta \lambda \rho Q_9 + k \zeta \eta \lambda \rho Q_{10} + \gamma \eta \lambda \rho^2 Q_{11}) \end{aligned}$$

$$\begin{aligned} \hat{\bar{\alpha}}_2 = & (\tilde{m}_o^2 \gamma \eta \lambda \bar{Q}_0 + \tilde{m}_o \bar{\zeta} \gamma \zeta \lambda \bar{Q}_1 + \bar{\gamma} \tilde{m}_o \zeta^2 \lambda \bar{Q}_2 + k \bar{\zeta} \zeta^2 \lambda \bar{Q}_3 + \bar{\zeta}^2 \gamma \zeta^2 \lambda \bar{Q}_4 + \bar{\gamma} \bar{\zeta} \zeta^3 \lambda \bar{Q}_5 \\ & + k \tilde{m}_o \zeta \eta \lambda \bar{Q}_6 + \bar{\zeta} \gamma \zeta \lambda \rho \bar{Q}_7 + \bar{\gamma} \zeta^2 \lambda \rho \bar{Q}_8 + \tilde{m}_o \gamma \eta \lambda \rho \bar{Q}_9 + k \zeta \eta \lambda \rho \bar{Q}_{10} + \gamma \eta \lambda \rho^2 \bar{Q}_{11}) \end{aligned}$$

$$\begin{aligned} \hat{\alpha}_3 = & (\bar{\gamma} \tilde{m}_o^2 \eta \lambda R_0 + \tilde{m}_o \bar{\zeta}^2 \gamma \lambda R_1 + \bar{\gamma} \tilde{m}_o \bar{\zeta} \zeta \lambda R_2 + k \bar{\zeta}^2 \zeta \lambda R_3 + \bar{\zeta}^3 \gamma \zeta \lambda R_4 + \bar{\gamma} \bar{\zeta}^2 \zeta^2 \lambda R_5 \\ & + k \tilde{m}_o \bar{\zeta} \eta \lambda R_6 + \bar{\zeta}^2 \gamma \lambda \rho R_7 + \bar{\gamma} \bar{\zeta} \zeta \lambda \rho R_8 + \bar{\gamma} \tilde{m}_o \eta \lambda \rho R_9 + k \bar{\zeta} \eta \lambda \rho R_{10} + \bar{\gamma} \eta \rho^2 \lambda R_{11}) \end{aligned}$$

$$\begin{aligned} \hat{\bar{\alpha}}_3 = & (\bar{\gamma} \tilde{m}_o^2 \eta \lambda \bar{R}_0 + \tilde{m}_o \bar{\zeta}^2 \gamma \lambda \bar{R}_1 + \bar{\gamma} \tilde{m}_o \bar{\zeta} \zeta \lambda \bar{R}_2 + k \bar{\zeta}^2 \zeta \lambda \bar{R}_3 + \bar{\zeta}^3 \gamma \zeta \lambda \bar{R}_4 + \bar{\gamma} \bar{\zeta}^2 \zeta^2 \lambda \bar{R}_5 \\ & + k \tilde{m}_o \bar{\zeta} \eta \lambda \bar{R}_6 + \bar{\zeta}^2 \gamma \lambda \rho \bar{R}_7 + \bar{\gamma} \bar{\zeta} \zeta \lambda \rho \bar{R}_8 + \bar{\gamma} \tilde{m}_o \eta \lambda \rho \bar{R}_9 + k \bar{\zeta} \eta \lambda \rho \bar{R}_{10} + \bar{\gamma} \eta \rho^2 \lambda \bar{R}_{11}) \end{aligned}$$

$$\begin{aligned}\hat{\alpha}_4 = & \sqrt{\frac{\lambda}{\eta}}(\tilde{m}_o^2 \gamma \eta^2 S_0 + \bar{\zeta}^2 \gamma \zeta^2 S_1 + \bar{\gamma} \bar{\zeta} \zeta^3 S_2 + \tilde{m}_o \bar{\zeta} \gamma \zeta \eta S_3 + \bar{\gamma} \tilde{m}_o \zeta^2 \eta S_4 + k \bar{\zeta} \zeta^2 \eta S_5 \\ & + k \tilde{m}_o \zeta \eta^2 S_6 + \bar{\zeta} \gamma \zeta \eta \rho S_7 + \bar{\gamma} \zeta^2 \eta \rho S_8 + \tilde{m}_o \gamma \eta^2 \rho S_9 + k \zeta \eta^2 \rho S_{10} + \gamma \eta^2 \rho^2 S_{11})\end{aligned}$$

$$\begin{aligned}\hat{\hat{\alpha}}_4 = & \sqrt{\frac{\lambda}{\eta}}(\bar{\gamma} \tilde{m}_o^2 \eta^2 \bar{S}_0 + \bar{\zeta}^3 \gamma \zeta \bar{S}_1 + \bar{\gamma} \bar{\zeta}^2 \zeta^2 \bar{S}_2 + \bar{\gamma} \tilde{m}_o \bar{\zeta} \zeta \eta \bar{S}_3 + \tilde{m}_o \bar{\zeta}^2 \gamma \eta \bar{S}_4 + k \bar{\zeta}^2 \zeta \eta \bar{S}_5 \\ & + k \tilde{m}_o \bar{\zeta} \eta^2 \bar{S}_6 + \bar{\gamma} \bar{\zeta} \zeta \eta \rho \bar{S}_7 + \bar{\zeta}^2 \gamma \eta \rho \bar{S}_8 + \bar{\gamma} \tilde{m}_o \eta^2 \rho \bar{S}_9 + k \bar{\zeta} \eta^2 \rho \bar{S}_{10} + \bar{\gamma} \eta^2 \rho^2 \bar{S}_{11})\end{aligned}$$

$$\begin{aligned}\hat{\alpha}_5 = & (\bar{\zeta}^2 \gamma \zeta \lambda T_0 + \bar{\gamma} \bar{\zeta} \zeta^2 \lambda T_1 + \tilde{m}_o \bar{\zeta} \gamma \eta \lambda T_2 + \bar{\gamma} \tilde{m}_o \zeta \eta \lambda T_3 + k \bar{\zeta} \zeta \eta \lambda T_4 \\ & + k \tilde{m}_o \eta^2 \lambda T_5 + \bar{\zeta} \gamma \eta \lambda \rho T_6 + \bar{\gamma} \zeta \eta \lambda \rho T_7 + k \eta^2 \lambda \rho T_8)\end{aligned}$$

$$\begin{aligned}\hat{\hat{\alpha}}_5 = & (\bar{\zeta}^2 \gamma \zeta \lambda \bar{T}_0 + \bar{\gamma} \bar{\zeta} \zeta^2 \lambda \bar{T}_1 + \tilde{m}_o \bar{\zeta} \gamma \eta \lambda \bar{T}_2 + \bar{\gamma} \tilde{m}_o \zeta \eta \lambda \bar{T}_3 + k \bar{\zeta} \zeta \eta \lambda \bar{T}_4 \\ & + k \tilde{m}_o \eta^2 \lambda \bar{T}_5 + \bar{\zeta} \gamma \eta \lambda \rho \bar{T}_6 + \bar{\gamma} \zeta \eta \lambda \rho \bar{T}_7 + k \eta^2 \lambda \rho \bar{T}_8)\end{aligned}$$

$$\begin{aligned}\hat{\alpha}_6 = & (\bar{\zeta}^2 \gamma \zeta \lambda U_0 + \bar{\gamma} \bar{\zeta} \zeta^2 \lambda U_1 + \tilde{m}_o \bar{\zeta} \gamma \eta \lambda U_2 + \bar{\gamma} \tilde{m}_o \zeta \eta \lambda U_3 + k \bar{\zeta} \zeta \eta \lambda U_4 + k \tilde{m}_o \eta^2 \lambda U_5 \\ & + \bar{\zeta} \gamma \eta \lambda \rho U_6 + \bar{\gamma} \zeta \eta \lambda \rho U_7 + k \eta^2 \lambda \rho U_8)\end{aligned}$$

$$\begin{aligned}\hat{\hat{\alpha}}_6 = & (\bar{\zeta}^2 \gamma \zeta \lambda \bar{U}_0 + \bar{\gamma} \bar{\zeta} \zeta^2 \lambda \bar{U}_1 + \tilde{m}_o \bar{\zeta} \gamma \eta \lambda \bar{U}_2 + \bar{\gamma} \tilde{m}_o \zeta \eta \lambda \bar{U}_3 + k \bar{\zeta} \zeta \eta \lambda \bar{U}_4 + k \tilde{m}_o \eta^2 \lambda \bar{U}_5 \\ & + \bar{\zeta} \gamma \eta \lambda \rho \bar{U}_6 + \bar{\gamma} \zeta \eta \lambda \rho \bar{U}_7 + k \eta^2 \lambda \rho \bar{U}_8)\end{aligned}$$

$$\begin{aligned}t_{(1,1)} &= -1 + x \\ t_{(2,1)} &= -3 + 4x + 3x^2 \\ t_{(2,2)} &= -1 + 5x^2 \\ t_{(2,3)} &= -1 + 2x + x^2 \\ t_{(2,4)} &= 1 - 6x + 7x^2 \\ t_{(2,5)} &= 1 - 5x + 2x^2 \\ t_{(2,6)} &= -3 + 3x + 2x^2 \\ t_{(3,1)} &= -3 + x + 5x^2 + 3x^3 \\ t_{(3,2)} &= (-1 + 3x)(1 + x^2) \\ t_{(3,3)} &= 2 - 9x + 6x^2 + 5x^3 \\ t_{(3,4)} &= -2 + 9x - 9x^2 + 4x^3\end{aligned}$$

$$\begin{aligned}
t_{(4,1)} &= 3 - 16x + 21x^2 - 6x^3 + 2x^4 \\
t_{(4,2)} &= 1 - x - 5x^2 + 5x^3 + 4x^4 \\
t_{(4,3)} &= 2 - 7x - 3x^2 + 11x^3 + 13x^4 \\
t_{(4,4)} &= 1 - 5x + 6x^2 - 5x^3 + 7x^4 \\
t_{(5,1)} &= -1 + 9x - 25x^2 + 29x^3 - 18x^4 + 14x^5 \\
t_{(5,2)} &= -2 + 8x - 3x^2 - 2x^3 - 8x^4 + x^5 \\
t_{(5,3)} &= 3 - 3x - 15x^2 - x^3 + 30x^4 + 14x^5 \\
t_{(5,4)} &= -1 - 3x + 11x^2 + 8x^3 - 12x^4 + 5x^5 \\
t_{(5,5)} &= -3 + 9x + 5x^2 - 10x^3 - 14x^4 + x^5 \\
t_{(6,1)} &= 1 + 4x - 38x^2 + 47x^3 + 10x^4 - 23x^5 + 7x^6 \\
t_{(7,1)} &= 1 - 8x + 25x^2 - 44x^3 + 64x^4 - 74x^5 + 36x^6 + 12x^7 \\
t_{(7,2)} &= 3 - 21x + 60x^2 - 114x^3 + 181x^4 - 147x^5 + 4x^6 + 18x^7 \\
t_{(7,3)} &= -1 + 5x - 3x^2 - 2x^3 - 39x^4 + 69x^5 - 37x^6 + 40x^7 \\
t_{(7,4)} &= 3 - 22x + 61x^2 - 102x^3 + 157x^4 - 142x^5 + 11x^6 + 18x^7 \\
t_{(7,5)} &= 2 + 11x - 72x^2 + 34x^3 + 123x^4 - 30x^5 - 49x^6 + 13x^7 \\
t_{(8,1)} &= 2 - 35x + 183x^2 - 398x^3 + 474x^4 - 441x^5 + 387x^6 - 170x^7 + 54x^8 \\
t_{(8,2)} &= 1 - 5x + 9x^2 - 10x^3 - 33x^4 + 195x^5 - 245x^6 + 44x^7 + 12x^8 \\
t_{(8,3)} &= 3 - 77x + 528x^2 - 1660x^3 + 2967x^4 - 3417x^5 + 2702x^6 - 1342x^7 + 360x^8 \\
t_{(8,4)} &= -1 + 19x - 115x^2 + 286x^3 - 211x^4 - 341x^5 + 767x^6 - 548x^7 + 128x^8 \\
t_{(8,5)} &= -3 + 23x - 81x^2 + 191x^3 - 307x^4 + 261x^5 - 91x^6 + 21x^7 + 18x^8 \\
t_{(9,1)} &= -1 + 12x - 39x^2 + 30x^3 + 79x^4 - 172x^5 + 63x^6 + 50x^7 - 6x^8 + 48x^9 \\
t_{(9,2)} &= 1 - 16x + 81x^2 - 174x^3 + 118x^4 + 222x^5 - 529x^6 + 500x^7 - 303x^8 + 84x^9 \\
t_{(9,3)} &= -1 + 21x - 135x^2 + 405x^3 - 599x^4 + 337x^5 + 127x^6 - 173x^7 - 128x^8 + 210x^9 \\
t_{(9,4)} &= 1 - 21x + 97x^2 - 155x^3 + 87x^4 - 29x^5 + 3x^6 + 63x^7 - 108x^8 + 30x^9 \\
t_{(10,1)} &= 5 - 51x + 219x^2 - 537x^3 + 883x^4 - 1025x^5 + 757x^6 - 287x^7 + 16x^8 + 12x^9 + 72x^{10} \\
t_{(10,2)} &= -1 + 21x - 152x^2 + 572x^3 - 1335x^4 + 2176x^5 - 2631x^6 + 2340x^7 \\
&\quad - 1380x^8 + 315x^9 + 171x^{10} \\
t_{(10,3)} &= -3 + 28x - 150x^2 + 486x^3 - 810x^4 + 422x^5 + 428x^6 - 670x^7 \\
&\quad + 605x^8 - 394x^9 + 186x^{10} \\
t_{(10,4)} &= 1 - 16x + 96x^2 - 338x^3 + 864x^4 - 1546x^5 + 1426x^6 + 114x^7 \\
&\quad - 1377x^8 + 762x^9 + 270x^{10} \\
t_{(11,1)} &= -1 + 18x - 121x^2 + 482x^3 - 1294x^4 + 1958x^5 - 226x^6 - 4462x^7 + 7239x^8 \\
&\quad - 4456x^9 + 483x^{10} + 252x^{11} \\
t_{(12,1)} &= 3 - 56x + 383x^2 - 1302x^3 + 2460x^4 - 2908x^5 + 2826x^6 - 2040x^7
\end{aligned}$$

$$-2121 x^8 + 8532 x^9 - 9449 x^{10} + 2766 x^{11} + 1674 x^{12}$$

$$\begin{aligned} p_2 &= (-1 + 2x)(1 + x) \\ p_3 &= -1 + 10x - 17x^2 + 12x^3 \\ p_4 &= (-1 + 3x)(-1 + 5x + x^3) \\ p_5 &= 1 - 7x + 21x^2 - 32x^3 + 20x^4 + 9x^5 \end{aligned}$$

$$\begin{aligned} P_0 &= -12 p_3 p_5 t_{(1,1)}^4 & P_1 &= 24 x^3 t_{(1,1)} t_{(10,1)} & P_2 &= -24 x t_{(10,2)} t_{(1,1)}^2 \\ P_3 &= 4 x t_{(11,1)} t_{(1,1)}^2 & P_4 &= 8 p_3 p_5 t_{(2,3)} t_{(1,1)}^3 & P_5 &= 4 x^2 p_3 p_5 t_{(1,1)}^4 \end{aligned}$$

$$\hat{\alpha}_1 = (\tilde{m}_o^2 \eta^2 \lambda P_0 + \bar{\zeta}^2 \zeta^2 \lambda P_1 + \tilde{m}_o \bar{\zeta} \zeta \eta \lambda P_2 + \bar{\zeta} \zeta \eta \lambda \rho P_3 + \tilde{m}_o \eta^2 \lambda \rho P_4 + \eta^2 \lambda \rho^2 P_5)$$

$$\begin{aligned} Q_0 &= 6\sqrt{3} p_3 p_4 t_{(1,1)}^5 & Q_1 &= 6\sqrt{3} x (-1 + 3x) t_{(8,1)} t_{(1,1)}^3 \\ Q_2 &= 6\sqrt{3} x^2 (-1 + 3x) p_2^2 t_{(3,1)} t_{(1,1)}^3 & Q_3 &= -12\sqrt{3} x^3 (-1 + 3x) t_{(1,1)} t_{(9,1)} \\ Q_4 &= \frac{-6\sqrt{3}}{\eta} x^3 (-3 + 5x) t_{(3,2)} t_{(4,1)} t_{(1,1)}^2 & Q_5 &= \frac{-6\sqrt{3}}{\eta} x^3 (-3 + 5x) p_2^2 t_{(1,1)}^2 t_{(3,2)} \\ Q_6 &= -12\sqrt{3} x^2 (-1 + 3x) p_3 t_{(4,2)} t_{(1,1)}^3 & Q_7 &= -2\sqrt{3} x (-1 + 3x) t_{(9,2)} t_{(1,1)}^3 \\ Q_8 &= -6\sqrt{3} x^3 (-1 + 3x) p_2^2 t_{(2,6)} t_{(1,1)}^4 & Q_9 &= -4\sqrt{3} p_3 p_4 t_{(2,3)} t_{(1,1)}^4 \\ Q_{10} &= 4\sqrt{3} x^3 (-1 + 3x) p_3 t_{(1,1)}^2 t_{(5,3)} & Q_{11} &= -2\sqrt{3} x^2 p_3 p_4 t_{(1,1)}^5 \end{aligned}$$

$$\begin{aligned} \bar{Q}_0 &= 6\sqrt{3} p_2 p_5 t_{(1,1)}^5 & \bar{Q}_1 &= 12\sqrt{3} x p_2 t_{(7,1)} t_{(1,1)}^3 \\ \bar{Q}_2 &= -12\sqrt{3} x^2 (-1 + 3x) p_2 t_{(5,2)} t_{(1,1)}^3 & \bar{Q}_3 &= 12\sqrt{3} x^3 p_2 t_{(7,2)} t_{(1,1)}^2 \\ \bar{Q}_4 &= \frac{-6\sqrt{3}}{\eta} x^3 t_{(3,2)} p_2^3 t_{(1,1)}^2 & \bar{Q}_5 &= \frac{-6\sqrt{3}}{\eta} x^3 p_2 t_{(3,2)} t_{(4,1)} t_{(1,1)}^2 \\ \bar{Q}_6 &= 24\sqrt{3} x^2 p_2 p_5 t_{(1,1)}^4 & \bar{Q}_7 &= -2\sqrt{3} x p_2 t_{(8,2)} t_{(1,1)}^3 \\ \bar{Q}_8 &= 2\sqrt{3} x^2 (-1 + 3x) p_2 t_{(6,1)} t_{(1,1)}^3 & \bar{Q}_9 &= -4\sqrt{3} p_2 p_5 t_{(2,3)} t_{(1,1)}^4 \\ \bar{Q}_{10} &= -4\sqrt{3} x^2 (1 + x) p_2 p_5 t_{(2,4)} t_{(1,1)}^2 & \bar{Q}_{11} &= -2\sqrt{3} x^2 p_2 p_5 t_{(1,1)}^5 \end{aligned}$$



$$\begin{aligned}
R_0 &= 6\sqrt{3}p_2p_5t_{(1,1)}^5 & R_1 &= -12\sqrt{3}x^2(-1+3x)p_2t_{(5,2)}t_{(1,1)}^3 \\
R_2 &= 12\sqrt{3}xp_2t_{(7,1)}t_{(1,1)}^3 & R_3 &= 12\sqrt{3}x^3p_2t_{(7,2)}t_{(1,1)}^2 \\
R_4 &= \frac{-6\sqrt{3}}{\eta}x^3p_2t_{(3,2)}t_{(4,1)}t_{(1,1)}^2 & R_5 &= \frac{-6\sqrt{3}}{\eta}x^3p_2^3t_{(3,2)}t_{(1,1)}^2 \\
R_6 &= 24\sqrt{3}x^2p_2p_5t_{(1,1)}^4 & R_7 &= 2\sqrt{3}x^2(-1+3x)p_2t_{(6,1)}t_{(1,1)}^3 \\
R_8 &= -2\sqrt{3}xp_2t_{(8,2)}t_{(1,1)}^3 & R_9 &= -4\sqrt{3}p_2p_5t_{(2,3)}t_{(1,1)}^4 \\
R_{10} &= -4\sqrt{3}x^2p_2p_5(1+x)t_{(2,4)}t_{(1,1)}^2 & R_{11} &= -2\sqrt{3}x^2p_2p_5t_{(1,1)}^5
\end{aligned}$$

$$\begin{aligned}
\bar{R}_0 &= 6\sqrt{3}p_3p_4t_{(1,1)}^5 & \bar{R}_1 &= 6\sqrt{3}x^2(-1+3x)p_2^2t_{(3,1)}t_{(1,1)}^3 \\
\bar{R}_2 &= 6\sqrt{3}x(-1+3x)t_{(8,1)}t_{(1,1)}^3 & \bar{R}_3 &= -12\sqrt{3}x^3(-1+3x)t_{(1,1)}t_{(9,1)} \\
\bar{R}_4 &= \frac{-6\sqrt{3}}{\eta}x^3(-3+5x)p_2^2t_{(1,1)}^2t_{(3,2)} & \bar{R}_5 &= \frac{-6\sqrt{3}}{\eta}x^3(-3+5x)t_{(1,1)}^2t_{(3,2)}t_{(4,1)} \\
\bar{R}_6 &= -12\sqrt{3}x^2(-1+3x)p_3t_{(4,2)}t_{(1,1)}^3 & \bar{R}_7 &= -6\sqrt{3}x^3(-1+3x)p_2^2t_{(2,6)}t_{(1,1)}^4 \\
\bar{R}_8 &= -2\sqrt{3}x(-1+3x)t_{(9,2)}t_{(1,1)}^3 & \bar{R}_9 &= -4\sqrt{3}p_3p_4t_{(2,3)}t_{(1,1)}^4 \\
\bar{R}_{10} &= 4\sqrt{3}x^3(-1+3x)p_3t_{(1,1)}^2t_{(5,3)} & \bar{R}_{11} &= -2\sqrt{3}x^2p_3p_4t_{(1,1)}^5
\end{aligned}$$

$$\begin{aligned}
S_0 &= -6\sqrt{2}p_3\hat{\sigma}t_{(3,4)}t_{(1,1)}^5 & S_1 &= -18\sqrt{2}x^3\hat{\sigma}t_{(4,1)}t_{(1,1)}^6 & S_2 &= -18\sqrt{2}\hat{\sigma}x^3p_2^2t_{(1,1)}^6 \\
S_3 &= -3\sqrt{2}x\hat{\sigma}t_{(8,3)}t_{(1,1)}^3 & S_4 &= -9\sqrt{2}x\hat{\sigma}p_2^2t_{(2,5)}t_{(1,1)}^5 & S_5 &= 12\sqrt{2}x^3\hat{\sigma}t_{(9,3)}t_{(1,1)} \\
S_6 &= 6\sqrt{2}x\hat{\sigma}p_3t_{(5,1)}t_{(1,1)}^3 & S_7 &= 3\sqrt{2}x\hat{\sigma}t_{(8,4)}t_{(1,1)}^4 & S_8 &= 3\sqrt{2}x\hat{\sigma}p_2^2t_{(4,4)}t_{(1,1)}^4 \\
S_9 &= 4\sqrt{2}p_3\hat{\sigma}t_{(2,3)}t_{(3,4)}t_{(1,1)}^4 & S_{10} &= -2\sqrt{2}x\hat{\sigma}p_3t_{(1,1)}^2t_{(7,3)} & S_{11} &= 2\sqrt{2}x^2\hat{\sigma}p_3t_{(3,4)}t_{(1,1)}^5
\end{aligned}$$

$$\begin{aligned}
\bar{S}_0 &= -6\sqrt{2}\hat{\sigma}p_3t_{(3,4)}t_{(1,1)}^5 & \bar{S}_1 &= -18\sqrt{2}x^3\hat{\sigma}p_2^2t_{(1,1)}^6 & \bar{S}_2 &= -18\sqrt{2}x^3\hat{\sigma}t_{(4,1)}t_{(1,1)}^6 \\
\bar{S}_3 &= -3\sqrt{2}x\hat{\sigma}t_{(8,3)}t_{(1,1)}^3 & \bar{S}_4 &= -9\sqrt{2}x\hat{\sigma}p_2^2t_{(2,5)}t_{(1,1)}^5 & \bar{S}_5 &= 12\sqrt{2}x^3\hat{\sigma}t_{(9,3)}t_{(1,1)} \\
\bar{S}_6 &= 6\sqrt{2}x\hat{\sigma}p_3t_{(5,1)}t_{(1,1)}^3 & \bar{S}_7 &= 3\sqrt{2}x\hat{\sigma}t_{(8,4)}t_{(1,1)}^4 & \bar{S}_8 &= 3\sqrt{2}x\hat{\sigma}p_2^2t_{(4,4)}t_{(1,1)}^4 \\
\bar{S}_9 &= 4\sqrt{2}\hat{\sigma}p_3t_{(2,3)}t_{(3,4)}t_{(1,1)}^4 & \bar{S}_{10} &= -2\sqrt{2}x\hat{\sigma}p_3t_{(1,1)}^2t_{(7,3)} & \bar{S}_{11} &= 2\sqrt{2}x^2\hat{\sigma}p_3t_{(3,4)}t_{(1,1)}^5
\end{aligned}$$

$$\begin{aligned}
T_0 &= 6x^2 (-1 + 3x) t_{(1,1)} t_{(10,3)} \\
T_2 &= 6x (-1 + 3x) p_3 t_{(5,4)} t_{(1,1)}^3 \\
T_4 &= -12x^2 t_{(12,1)} \\
T_6 &= -2x (-1 + 3x) p_3 t_{(1,1)}^2 t_{(7,5)} \\
T_8 &= 4x p_3 p_5 t_{(1,1)} t_{(4,3)}
\end{aligned}$$

$$\begin{aligned}
T_1 &= 18x^2 p_2 t_{(7,4)} t_{(1,1)}^3 \\
T_3 &= 18x p_2 p_5 t_{(1,1)}^5 \\
T_5 &= -12x p_3 p_5 t_{(1,1)}^2 t_{(2,2)} \\
T_7 &= -6x p_2 p_5 t_{(3,3)} t_{(1,1)}^3
\end{aligned}$$

$$\begin{aligned}
\bar{T}_0 &= 18x^2 p_2 t_{(7,4)} t_{(1,1)}^3 \\
\bar{T}_2 &= 18x p_2 p_5 t_{(1,1)}^5 \\
\bar{T}_4 &= -12x^2 t_{(12,1)} \\
\bar{T}_6 &= -6x p_2 p_5 t_{(3,3)} t_{(1,1)}^3 \\
\bar{T}_8 &= 4x p_3 p_5 t_{(1,1)} t_{(4,3)}
\end{aligned}$$

$$\begin{aligned}
\bar{T}_1 &= 6x^2 (-1 + 3x) t_{(1,1)} t_{(10,3)} \\
\bar{T}_3 &= 6x (-1 + 3x) p_3 t_{(5,4)} t_{(1,1)}^3 \\
\bar{T}_5 &= -12x p_3 p_5 t_{(2,2)} t_{(1,1)}^2 \\
\bar{T}_7 &= -2x (-1 + 3x) p_3 t_{(7,5)} t_{(1,1)}^2
\end{aligned}$$

$$\begin{aligned}
U_0 &= 6\sqrt{3}i x^2 (-1 + 3x) t_{(9,4)} t_{(1,1)}^2 \\
U_2 &= -6\sqrt{3}i x (-1 + 3x) p_3 t_{(5,5)} t_{(1,1)}^3 \\
U_4 &= -12\sqrt{3}i x^2 t_{(10,4)} t_{(1,1)}^2 \\
U_6 &= 2\sqrt{3}i x^2 (-1 + 3x) p_3 t_{(5,4)} t_{(1,1)}^3 \\
U_8 &= -4\sqrt{3}i x^2 p_3 p_5 t_{(2,2)} t_{(1,1)}^2
\end{aligned}$$

$$\begin{aligned}
U_1 &= 6\sqrt{3}i x^2 p_2 t_{(8,5)} t_{(1,1)}^2 \\
U_3 &= -6\sqrt{3}i x p_2 p_5 t_{(2,1)} t_{(1,1)}^3 \\
U_5 &= 12\sqrt{3}i x p_3 p_5 t_{(1,1)}^4 \\
U_7 &= 6\sqrt{3}i x^2 p_2 p_5 t_{(1,1)}^5
\end{aligned}$$

$$\begin{aligned}
\bar{U}_0 &= -6\sqrt{3}i x^2 p_2 t_{(8,5)} t_{(1,1)}^2 \\
\bar{U}_2 &= 6\sqrt{3}i x p_2 p_5 t_{(2,1)} t_{(1,1)}^3 \\
\bar{U}_4 &= 12\sqrt{3}i x^2 t_{(10,4)} t_{(1,1)}^2 \\
\bar{U}_6 &= -6\sqrt{3}i x^2 p_2 p_5 t_{(1,1)}^5 \\
\bar{U}_8 &= 4\sqrt{3}i x^2 p_3 p_5 t_{(2,2)} t_{(1,1)}^2
\end{aligned}$$

$$\begin{aligned}
\bar{U}_1 &= -6\sqrt{3}i x^2 (-1 + 3x) t_{(9,4)} t_{(1,1)}^2 \\
\bar{U}_3 &= 6\sqrt{3}i x (-1 + 3x) p_3 t_{(1,1)}^3 t_{(5,5)} \\
\bar{U}_5 &= -12\sqrt{3}i x p_3 p_5 t_{(1,1)}^4 \\
\bar{U}_7 &= -2\sqrt{3}i x^2 (-1 + 3x) p_3 t_{(5,4)} t_{(1,1)}^3
\end{aligned}$$

## References

- [1] C. S. Aulakh, K. Benakli and G. Senjanovic, Phys. Rev. Lett. **79**, 2188 (1997) [arXiv:hep-ph/9703434]; C. S. Aulakh, A. Melfo and G. Senjanovic, Phys. Rev.

- D **57** (1998) 4174 [arXiv:hep-ph/9707256]; C. S. Aulakh, A. Melfo, A. Rasin and G. Senjanovic, Phys. Lett. B **459**, 557 (1999) [arXiv:hep-ph/9902409].
- [2] C. S. Aulakh, B. Bajc, A. Melfo, A. Rasin and G. Senjanovic, Nucl. Phys. B **597**, 89 (2001) [arXiv:hep-ph/0004031].
- [3] C.S.Aulakh and A. Girdhar, hep-ph/0204097; v2 August 2003; v4, 9 February, 2004; Int. J. Mod. Phys. A **20**, 865 (2005)
- [4] B. Bajc, A. Melfo, G. Senjanovic and F. Vissani, Phys. Rev. D **70**, 035007 (2004) [arXiv:hep-ph/0402122].
- [5] C. S. Aulakh and A. Girdhar, Nucl. Phys. B **711**, 275 (2005).
- [6] T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac and N. Okada, Eur. Phys. J. C **42**, 191 (2005) arXiv:hep-ph/0401213v1,v2.
- [7] K. Y. Oda, E. Takasugi, M. Tanaka and M. Yoshimura, Phys. Rev. D **59**, 055001 (1999) [arXiv:hep-ph/9808241]; K. Matsuda, Y. Koide, T. Fukuyama and H. Nishiura, Phys. Rev. D **65**, 033008 (2002) [Erratum-ibid. D **65**, 079904 (2002)] [arXiv:hep-ph/0108202] ; K. Matsuda, Y. Koide and T. Fukuyama, Phys. Rev. D **64**, 053015 (2001) [arXiv:hep-ph/0010026]. N. Oshimo, ; Phys. Rev. D **66**, 095010 (2002) [arXiv:hep-ph/0206239]; N. Oshimo, Nucl. Phys. B **668**, 258 (2003) [arXiv:hep-ph/0305166]; B. Bajc, G. Senjanovic and F. Vissani, Phys. Rev. Lett. **90** (2003) 051802 [arXiv:hep-ph/0210207]; H. S. Goh, R. N. Mohapatra and S. P. Ng, Phys. Lett. B **570**, 215 (2003) [arXiv:hep-ph/0303055]. H. S. Goh, R. N. Mohapatra and S. P. Ng, Phys. Rev. D **68**, 115008 (2003) [arXiv:hep-ph/0308197]. H. S. Goh, R. N. Mohapatra and S. Nasri, Phys. Rev. D **70** (2004) 075022 [arXiv:hep-ph/0408139]; S. Bertolini, M. Frigerio and M. Malinsky, Phys. Rev. D **70**, 095002 (2004) [arXiv:hep-ph/0406117]; S. Bertolini and M. Malinsky, arXiv:hep-ph/0504241 ; K. S. Babu and C. Macesanu, arXiv:hep-ph/0505200. B. Bajc, G. Senjanovic and F. Vissani, Phys. Rev. D **70**, 093002 (2004) [arXiv:hep-ph/0402140]; B. Bajc, G. Senjanovic and F. Vissani, arXiv:hep-ph/0110310; K. S. Babu and C. Macesanu, Phys. Rev. D **72**, 115003 (2005) [arXiv:hep-ph/0505200].
- [8] P. Minkowski, Phys. Lett. **B67**,110(1977); M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, eds. P. van Nieuwenhuizen and D.Z. Freedman (North Holland 1979); T. Yanagida, in *Proceedings of Workshop on Unified Theory and Baryon number in the Universe*, eds. O. Sawada and A. Sugamoto (KEK 1979); R.N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980); R.N. Mohapatra and G. Senjanović, Phys. Rev. **D23**,165 (1981); G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. **B181**, 287 (1981).

- [9] K.S.Babu and R.N.Mohapatra, Phys. Rev. Lett. **70**(1993)2845.
- [10] C.S. Aulakh and R.N. Mohapatra, CCNY-HEP-82-4 April 1982, CCNY-HEP-82-4-REV, Jun 1982 , Phys. Rev. **D28**, 217 (1983).
- [11] T.E. Clark, T.K.Kuo, and N.Nakagawa, Phys. Lett. **115B**, 26(1982).
- [12] C. S. Aulakh, B. Bajc, A. Melfo, G. Senjanovic and F. Vissani, Phys. Lett. B **588**, 196 (2004) [arXiv:hep-ph/0306242].
- [13] R. N. Mohapatra, Phys. Rev. D **34**, 3457 (1986).; A. Font, L. E. Ibanez and F. Quevedo, Phys. Lett. B **228**, 79 (1989); S. P. Martin, Phys. Rev. D **46**, 2769 (1992) [arXiv:hep-ph/9207218]; D. G. Lee and R. N. Mohapatra, Phys. Rev. D **51**, 1353 (1995) [arXiv:hep-ph/9406328].
- [14] C. S. Aulakh, *From germ to bloom*, arXiv:hep-ph/0506291.
- [15] C. S. Aulakh and S. K. Garg, Nucl. Phys. B **757**, 47 (2006) [arXiv:hep-ph/0512224].
- [16] C. S. Aulakh, *Fermion mass hierarchy in the Nu MSGUT. I: The real core*, arXiv:hep-ph/0602132 ;
- [17] C. S. Aulakh, *MSGUT Reborn ?* arXiv:hep-ph/0607252
- [18] W. Grimus and H. K  hnb  ck, *Fermion masses and mixings in a renormalizable  $SO(10) \times Z_2$  GUT* , hep-ph/0607197.
- [19] B. Bajc, A. Melfo, G. Senjanovic and F. Vissani, AIP Conf. Proc. **805**, 152 (2006) [AIP Conf. Proc. **805**, 326 (2006)] [arXiv:hep-ph/0511352].
- [20] C. S. Aulakh, Phys. Rev. D **72**, 051702 (2005), arXiv:hep-ph/0501025.
- [21] H. Murayama and A. Pierce, Phys. Rev. D **65**, 055009 (2002), [arXiv:hep-ph/0108104].
- [22] C. S. Aulakh, “*Truly minimal unification: Asymptotically strong panacea?*,” arXiv:hep-ph/0207150.
- [23] B. Bajc and G. Senjanovic, “*Proton decay, supersymmetry breaking and its mediation,*” arXiv:hep-ph/0611308.
- [24] B. A. Ovrut and S. Raby, Phys. Lett. B **138** (1984) 72 and Phys. Rev. D **31** (1985) 2968; M. Drees, Phys. Rev. D **33** (1986) 1468; K. Agashe, Phys. Lett. B **444** (1998) 61 [arXiv:hep-ph/9809421] and Nucl. Phys. B **588** (2000) 39 [arXiv:hep-ph/0003236] ; F. R. Joaquim and A. Rossi, arXiv:hep-ph/0604083 and arXiv:hep-ph/0607298; R. Dermisek, H. D. Kim and I. W. Kim, arXiv:hep-ph/0607169;.

- [25] C. S. Aulakh, “*Taming asymptotic strength*,” arXiv:hep-ph/0210337.
- [26] N. Seiberg, Phys. Lett. **B318**, 469(1993); N. Seiberg, Phys. Rev.**D49**, 6857 (1994). For reviews and complete references see : K.Intriligator, N.Seiberg, Proc. of ICTP Summer School, 1995, Nucl. Phys. Proc. Suppl. **45BC**,1(1996), hep-th/9509066; M.Shifman,Lectures at ICTP Summer School, 1996, Prog. Part. Nucl. Phys. **39**,1(1997).
- [27] F. David, “A Comment On Induced Gravity,” Phys. Lett. B **138**, 383 (1984); F. David and A. Strominger, “On The Calculability Of Newton’s Constant And The Renormalizability Of Scale Invariant Quantum Gravity,” Phys. Lett. B **143** (1984) 125.
- [28] N. Arkani-Hamed, M. Porrati and L. Randall, JHEP **0108** (2001) 017 [arXiv:hep-th/0012148].
- [29] A. Melfo and G. Senjanovic, “*Supersymmetric grand unification: The quest for the theory*,” arXiv:hep-ph/0511011.
- [30] E. Cremmer, S. Ferrara, L. Girardello and A. Van Proeyen, Nucl. Phys. B **212**, 413 (1983); Phys. Lett. B **116**, 231 (1982). E. Cremmer, B. Julia, J. Scherk, S. Ferrara, L. Girardello and P. van Nieuwenhuizen,
- [31] T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac and N. Okada, J. Math. Phys. **46** (2005) 033505 [arXiv:hep-ph/0405300].
- [32] L.J.Hall, Nucl. Phys. **B178**,75(1981).
- [33] M. Drees, R.M. Godbole and P.Roy, Theory and Phenomenology of Sparticles, World Scientific, 2004. R. Barbieri,
- [34] L. J. Hall and A. Strumia, Nucl. Phys. B **445** (1995) 219 [arXiv:hep-ph/9501334]; R. Barbieri, L. J. Hall and A. Strumia, Nucl. Phys. B **445** (1995) 219 [arXiv:hep-ph/9501334].
- [35] C. S. Aulakh and S. K. Garg, to appear.